

## СПИСЪК

на забелязаните цитирания на научните публикации

на доц. д-р Светлана Петкова Мишева (бивша фамилия Ланджева)

**Общ брой цитирания: 464 (към 24.11.2017)**

**От тях:**

- в списания с IF: 230 (вкл. 21 с RG IF)
- в дисертации в България: 12
- в дисертации в чужбина: 48
- в книги: 25

**Landjeva S, Todorova M, Ganeva G (1994) Immunogenetic investigation on the leaf rust (*Puccinia recondita* Rob. ex Desm. f.sp. *tritici* Erikss.) resistance in tetraploid wheats, Раст. науки 31 (7-10): 266-270**

1. Кържин Хр (2003) Проучвания върху ръждите по пшеницата в България и средствата за борба с тях. Добруджански Земеделски Институт, Генерал Тошево, стр. 128

**Landgeva S, Ganeva G, Georgieva V (1995) Somatic karyotypes and N-banding patterns of tetraploid wheats. Biol Zentralblatt 114: 253-265**

2. Belay G, Merker A (1998) Cytogenetic analysis of a spontaneous 5B/6B translocation in tetraploid wheat landraces from Ethiopia, and implications for breeding. Plant Breeding 117: 37-542 (IF 2016-1.335)
3. Belay G, Merker A (1999) C-band polymorphism and chromosomal rearrangements in tetraploid wheat (*Triticum turgidum* L.) landraces from Ethiopia. Wheat Information Service 88: 6-14
4. Vassileva I, Balevska P, Tsvetkov S (2003) Chromosome constitution of triticale cultivar Zaryad. Comptes rendus de l'Académie bulgare des Sciences 56 (1): 85-88 (IF 2016 – 0.251)

**Landjeva S, Ganeva G (1996) N-banded karyotype of *Aegilops ovata* L. and chromosomal constitution of its amphiploid with *Triticum aestivum* L., Plant Breed 115: 330-334**

5. Спецов П (1998) Използване на видове от *Aegilops* (2n=28, UUSS) за подобряване на устойчивостта към брашнеста мана и други стопански признания при обикновената зимна пшеница. Дисертация за присъждане на научната степен "Доктор на науките", СА, София
6. Пламенов Д (2003) Разпространение и характеристика на диви житни (*Aegilops* и *Triticum*) по Черноморието. Дисертация за присъждане на образователната и научна степен "Доктор", ТУ - Варна.
7. Szinay D (2010) The development of FISH tools for genetic, phylogenetic and breeding studies in tomato (*Solanum lycopersicum*). PhD Thesis, Wageningen University, Wageningen, NL, 147 pages

**Landjeva S, Todorova M, Ganeva G (1998) Development and cytogenetic characterization of a disomic wheat-*Aegilops ovata* chromosome addition line with leaf rust resistance. J Genet Breed (Rome) 52: 217-222**

8. Stoilova TB (2002) Biochemical diagnostics of *Aegilops geniculata* Roth chromosomes in homoeologous group 3 of wheat. Journal of Genetics and Breeding (Rome) 56 (4): 359-364
9. Кържин Хр (2003) Проучвания върху ръждите по пшеницата в България и средствата за борба с тях. Добруджански Земеделски Институт, Генерал Тошево, стр. 128

**Landjeva S, Merakchijkska M, Ganeva G (1998). Seedling copper tolerance and cytogenetic characterization of wheat-Ae. ovata hybrid lines. Wheat Inform Serv 87: 27-30**

10. Пламенов Д, Спецов П (2003) Видов състав и разпространение на дивите житни (*Aegilops* и *Triticum*) в България. Растениевъдни Науки 40 (3): 197-203
11. Пламенов Д (2003) Разпространение и характеристика на диви житни (*Aegilops* и *Triticum*) по Черноморието. Дисертация за присъждане на образователната и научна степен "Доктор", ТУ - Варна
12. Vassilev A, Lidon F, Scotti Campos P, Ramalho JC, Barreiro MG, Yordanov I (2003) Cu-induced changes in chloroplast lipids and photosystem 2 activity in barley plants. Bulgarian Journal of Plant Physiology 29 (1-2): 33-43
13. Bálint A, Kovacs G, Börner A, Galiba G, Sutka J (2003) Substitution analysis of seedling stage copper tolerance in wheat. Acta Agronomica Hungarica 51(4): 397-404
14. Bálint A, Röder MS, Hell R, Galiba G, Börner A (2007) Mapping of QTLs affecting copper tolerance and the Cu, Fe, Mn and Zn contents in the shoots of wheat seedlings. Biologia Plantarum 51:129-134 (IF 2016 – 1.551)

**Landjeva S, Ganeva G, Petrova N, Spetsov P (1998). Cytogenetical characterization of disease resistant wheat-Aegilops ovata derivatives. Proc 9th Int Wheat Genet Symp, v.2, Section 1, pp 64-66**

15. Пламенов Д (2003) Разпространение и характеристика на диви житни (*Aegilops* и *Triticum*) по Черноморието. Дисертация за присъждане на научната и образователна степен "Доктор", ТУ - Варна.

**Landjeva S, Ganeva G (1998). Transfer of *Aegilops ovata* chromosomes into bread wheat. EWAC Newsletter 10: 29-34**

16. Wojciechowska B, Pudelska H (2002) Hybrids and amphiploids of *Aegilops ovata* L. with *Secale cereale* L.: Production, morphology and fertility. Journal of Applied Genetics 43 (4): 415-421 (IF 2016 – 1.665)
17. Khan AA, Khaliq M, Nawaz-ur-Rehman MS, Irshad A (2005) Cytogenetics and Evolution of *Triticum aestivum* L. em Thell. International Journal of Agriculture and Biology 7(3): 527-534
18. Rehman M, Hansen JL, Zemetra RS (2006) Hybrids and amphiploids of *Aegilops cylindrica* with *Triticum aestivum* L.; production morphology and fertility. Pakistan Journal of Biological Sciences 9: 1563-1566 (RG IF 2016 – 0.73)
19. Loureiro I, Escorial C, García-Baudin JM, Chueca MC (2009) Spontaneous wheat-*Aegilops biuncialis*, *Ae. geniculata* and *Ae. triuncialis* amphiploid production, a potential way of gene transference. Spanish Journal of Agricultural Research 7: 614-620 (IF 2016 - 0.687)
20. Apolinarska B, Wioeniewska H, Wojciechowska B (2010) *Aegilops*-rye amphiploids and substitution rye used for introgression of genetic material into rye (*Secale cereale* L.) Journal of Applied Genetics 51:413–420 (IF 2016 – 1.665)

**Landjeva S, Ganeva G (1999). Identification of *Aegilops ovata* chromosomes added to the wheat (*Triticum aestivum* L.) genome. Cereal Res Comm 27: 55-61**

21. Wojciechowska B, Pudelska H (2002) Hybrids and amphiploids of *Aegilops ovata* L. with *Secale cereale* L.: Production, morphology and fertility. Journal of Applied Genetics 43:415-421 (IF 2016 – 1.665)
22. Singh Ram J (2002) Plant Cytogenetics, 2nd Ed. CRC Press, pp 512
23. Пламенов Д, Спецов П (2003) Видов състав и разпространение на дивите житни (*Aegilops* и *Triticum*) в България. Растениевъдни науки 40 (3): 197-203
24. Numerova OM, Pershina LA, Salina EA, Shumny VK (2004) Barley chromosome identification using genomic *in situ* hybridization in the genome of backcrossed progeny of barley-wheat amphiploids [*Hordeum geniculatum* All. (2n=28) x *Triticum aestivum* L. (2n=42)] (2n=70) Russian Journal of Genetics 40 (9): 1007-1010 (IF 2016 – 0.55)
25. Loureiro I, Escorial C, García-Baudin JM, Chueca MC (2009) Spontaneous wheat-*Aegilops biuncialis*, *Ae. geniculata* and *Ae. triuncialis* amphiploid production, a potential way of gene

transference. Spanish Journal of Agricultural Research 7:614-620 (IF 2016 – 0.687)

26. Apolinarska B, Wioeniewska H, Wojciechowska B (2010) *Aegilops*-rye amphiploids and substitution rye used for introgression of genetic material into rye (*Secale cereale* L.) Journal of Applied Genetics 51: 413–420 (IF 2016 – 1.665)
27. Kwiatek M, Błaszczyk L, Wiśniewska H and Apolinarska B (2012) *Aegilops-Secale* amphiploids: chromosome categorisation, pollen viability and identification of fungal disease resistance genes. Journal of Applied Genetics 53 (1): 37-40 (IF 2016 – 1.665)

**Landjeva S, Ganeva G (2000). Chromosome N-banding polymorphism in *Aegilops geniculata* Roth. Genetic Resources and Crop Evolution 47: 35-42**

28. Ekmekci Y, Terzioglu S (2005) Effects of oxidative stress induced by paraquat on wild and cultivated wheats. Pesticide Biochemistry and Physiology 83(2-3):69-81 (IF 2016 – 2.59)
29. Fletcher, R. Australian New Crops Newsletter. Listing of Interesting Plants of the World; <http://www.newcrops.uq.edu.au/>
30. Molnár I, Cifuentes M, Schneider A, Benavente E, Molnár-Láng M (2011) Association between simple sequence repeat-rich chromosome regions and intergenomic translocation breakpoints in natural populations of allopolyploid wild wheats. Annals of Botany 107: 65-76 (IF 2016 – 4.041)
31. Schneider A, Benavente E, Molnár-Láng M (2012) Detection of Various U and M chromosomes in wheat-*Aegilops biuncialis* Hybrids and Derivatives Using Fluorescence *in situ* hybridisation and molecular markers. Czech Journal of Genetics and Plant Breeding 48:169–177 (IF 2016 – 0.532)

**Ланджева С (2000). Хромозомен анализ на *Aegilops geniculata* Roth и негови хибриди с пшеницата (*Triticum aestivum* L.), проявяващи важни стопански признаки, Дисертация за присъждане на образователната и научна степен “доктор”, ИГ-БАН, София.**

32. Stoilova TB (2002) Biochemical diagnostics of *Aegilops geniculata* Roth chromosomes in homoeologous group 3 of wheat. Journal of Genetics and Breeding (Rome) 56 (4): 359-364
33. Пламенов Д, Спецов П (2003). Видов състав и разпространение на дивите житни (*Aegilops* и *Triticum*) в България. Растениевъдни Науки 40 (3): 197-203
34. Пламенов Д (2003). Разпространение и характеристика на диви житни (*Aegilops* и *Triticum*) по Черноморието. Дисертация за присъждане на образователната и научна степен “Доктор”, ТУ - Варна

**Ganeva G, Landjeva S (2003). Identification of chromosome translocations in the Bulgarian wheat variety ‘Gladiator 113’. Wheat Inform Serv 97: 23-24**

35. Schlegel R (2008) Current list of wheats with rye introgression. V05-08, 1-14, <http://www.desicca.de/Wheat-rye introgression>

**Ganeva G, Landjeva S (2004). Genome structure of Bulgarian bread wheat cultivars as revealed by meiotic chromosome pairing and interchange analysis. J Genet Breed (Rome) 58: 265-272**

36. Schlegel R (2004) A compendium of reciprocal translocation in wheat. PBU. [www.plant.breeding.privat.t-online.de](http://www.plant.breeding.privat.t-online.de)

**Landjeva S, Merakchijska-Nikolova M, Ganeva G (2003). Copper toxicity tolerance in *Aegilops* and *Haynaldia* seedlings. Biol Plantarum 46: 479-480**

37. Bernal M (2006) Mecanismos de tolerancia al exceso de cobre en suspensiones celulares de soja. Caracterización del transportador de cobre HMA8. PhD Thesis. Estación Experimental de Aula Dei; Universidad de Zaragoza
38. Groppa MD, Zawoznik MS, Tomaro ML, Benavides MP (2008) Inhibition of root growth and polyamine metabolism in sunflower (*Helianthus annuus*) seedlings under cadmium and copper stress. Biological Trace Element Research 126(1-3):246-256 (IF 2016 - 1.307)
39. Wang Yu-Hai, Wang Li-Ming, Bao Yin-Guang, Cui Fa, Hao Yuan-Feng, Zong Hao, LiXing-Feng, Gao

- Ju-Rong, and Wang Hong-Gang (2009) Segregations of chromosome and trait in hybrid generations derived from cross between *Triticum aestivum* and amphidiploid of *Aegilops ventricosa* × *Aegilops cylindrica*. *Acta Agronomica Sinica* 35:1261–1267 (RG IF 2016 – 0.12)
40. Molnár I and Molnár-Lág M (2010) GISH reveals different levels of meiotic pairing with wheat for individual *Aegilops biuncialis* chromosomes. *Biologia Plantarum* 54 (2): 259-264 (IF 2016 – 1.551)
  41. Stoyanov H (2013) Characteristics of second generation hybrids between wheat species (*Triticum* sp.) and *Aegilops cylindrica* Host. *Publications of the University of Agronomic Sciences and Veterinary Medicine of Bucharest: Scientific Papers. Series A. Agronomy*, Vol. LVI, pp 357-362
  42. Stoyanov H (2013) Characteristics of natural hybrids between *Aegilops cylindrica* HOST. and common winter wheat (*Triticum aestivum* L.). *AgroLife Scientific Journal* 2(1): 66-71
  43. Stoyanov H (2013) Status of remote hybrids in the *Poaceae*: problems and prospects. *Agricultural Science and Technology* 5(1), 3 – 12
  44. Dimitrijević M, Petrović S (2014). Species variation of *Aegilops* genus and heavy metal content in plant habitat soil at southern adriatic localities. *Genetika* 46 (2) 381-400 (RG IF 2016 – 0.44)
  45. Wang Yu-ha , HE Fang, Bao Yin-guang, Ming Dong-feng, Dong Lei, Han Qing-dian, Li Ying-ying, Wang Hong-gang (2016) Development and genetic analysis of a novel wheat-Aegilops germplasm TA002 resistant to powdery mildew. *Scientia Agricultura Sinica* 49(3): 418-428
  46. Baltrénaitė E, Baltrénas P, Lietuvninkas A (2016) Biogeochemical and Functional Traits of a Tree in Metal-Contaminated Territory. Chapter In: *The Sustainable Role of the Tree in Environmental Protection Technologies*, Springer, ISBN: 978-3-319-25475-3, pp 127-148
  47. Agami RA (2016) Pre-soaking in indole-3-acetic acid or spermidine enhances copper tolerance in wheat seedlings. *South African Journal of Botany* 104:167-174 (IF 2016 - 1.427)

**Ganeva G, Landjeva S, Merakchijkska M (2003/2004). Effects of chromosome substitutions on copper toxicity tolerance in wheat seedlings. Biol Plantarum 47: 621-623**

48. Bálint A, Röder MS, Hell R, Galiba G, Sutka J, Börner A (2006) Cereals with better heavy metal tolerance and nutritional value: physical and genetic mapping of copper tolerance and shoot micronutrient content in wheat. *EWAC Newsletter* 13: 55-58
49. Bálint A, Röder MS, Hell R, Galiba G, Börner A (2007) Mapping of QTLs affecting copper tolerance and the Cu, Fe, Mn and Zn concentrations in the shoots of wheat seedlings. *Biologia Plantarum* 51: 129-134 (IF 2016 – 1.551)
50. Arraiano LS, Kirby J, Brown JKM (2007) Cytogenetic analysis of the susceptibility of the wheat line Hobbit sib (Dwarf A) to *Septoria tritici* blotch. *Theoretical and Applied Genetics* 116: 113-122 (IF 2016 – 4.132)
51. Bálint A, Szira F, Röder MS, Galiba G, Börner A (2009) Mapping of loci affecting copper tolerance in wheat - The possible impact of the vernalization gene *Vrn-A1*. *Environmental and Experimental Botany* 65: 369-375 (IF 2016 - 4.369)
52. Mehes-Smith M, Nkongolo K, Cholewa E (2013) Coping Mechanisms of Plants to Metal Contaminated Soil. *Environmental Change and Sustainability*, Chapter 3, pp 53-90

**Landjeva S, Angelov G, Nenova V, Merakchijkska M and Ganeva G (2004). Seedling growth and peroxidase responses to excess copper in wheat-Aegilops geniculata chromosome addition and substitution lines. Genetics and Breeding 33 (3-4): 11-19**

53. Kosturkova G and Tasheva K (2009). Chapter 7. Women's Leadership in Plant Biotechnology and Related Sciences. In: Nardi V (Ed) *Feminism and Women in Leadership*, NOVA Publishers NY, pp 127-143

**Ganeva G, Korzun V, Landjeva S, Tsenov N, Atanasova M (2005). Identification, distribution and effects on agronomical traits of the semi-dwarfing Rht alleles in Bulgarian bread wheat varieties. Euphytica 145 (3):**

54. Zhang X, Yang S, Zhou Y, He Z, Xia X (2006) Distribution of the *Rht-B1b*, *Rht-D1b* and *Rht8* reduced height genes in autumn-sown Chinese wheats detected by molecular markers. *Euphytica* 152 (1): 109-116 (IF 2016 – 1.626)
55. Bognár Z, Láng L, Bedő Z (2007) Effect of environment on the plant height of wheat germplasm. *Cereal Research Communications* 35:281-284 (IF 2016 – 0.496)
56. Ellis MH, Bonnett DG, Rebetzke GJ (2007) A 192bp allele at the *Xgwm261* locus is not always associated with the *Rht8* dwarfing gene in wheat (*Triticum aestivum* L.) *Euphytica* 157 (1-2): 209-214 (IF 2016 – 1.626)
57. Tošovic-Maric B, Kobiljski B, Obreht D, Vapa Lj (2008) Evaluation of wheat *Rht* genes using molecular markers. *Genetika* 40 (1): 31-38 (RG IF 2016 – 0.44)
58. Zheleva D (2008). Дисертация за присъждане на образователната и научна степен "Доктор", Агробиоинститут, НЦАН, София
59. Žofajová A, Užík M, Muchová D, Lichvárová M, Ondrejčák F, Mihálik D (2008). The evaluation of tall and dwarf parental winter wheat cultivars. *Agriculture (Poľnohospodárstvo)*, 54 (4): 182–193
60. Al-Ajlouni ZI (2008) The effect of semi-dwarfing genes in wheat on plant height in diverse environment in Nebraska. PhD Thesis, The University of Nebraska – Linco
61. Šíp V, Chrpová J, Žofajová A, Panková K, Užík M and Snape JW (2009) Effects of specific *Rht* and *Ppd* alleles on agronomic traits in winter wheat cultivars grown in middle Europe. *Euphytica* 172(2): 221-233 (IF 2016 – 1.626)
62. Dvojković K, Šatović Z, Drezner G, Somers DJ, Lalić A, Novoselović D, Horvat D, Marić S, Šarčević H (2010). Allelic variability of croatian wheat cultivars at the microsatellite locus *Xgwm261*. *Poljoprivreda* 16 (1): 32-37
63. Kang S, Bai F, Lan S, Li X (2010) Research Progress of Wheat Dwarfing Genes. *Journal of Hebei Normal University (Natural Science Ed)* 34(1) DOI: CNKI: SUN:HBSZ.0.2010-01-025
64. Voss, H-H (2010) Inheritance of quantitative resistance and aggressiveness in the wheat / *Fusarium* pathosystem with emphasis on *Rht* dwarfing genes. PhD Thesis, Universität Hohenheim
65. Heyns IC (2010) Mapping and restructuring of an *Ae. kotschy* derived translocation segment in common wheat. PhD thesis University of Stellenbosch, South Africa
66. Šíp V, Chrpová J, Žofajová A, Milec Z, Mihalik D, Pánková, K, Snape JW (2011) Evidence of selective changes in winter wheat in middle-European environments reflected by allelic diversity at loci affecting plant height and photoperiodic response. *Journal of Agricultural Science* 149:313-326 (IF 2016 - 1.291)
67. Voss H-H, RL Bowden, JF Lesliej and T Miedaner (2010) Variation and transgression of aggressiveness among two *Gibberella zae* crosses developed from highly aggressive parental isolates. *Phytopathology* 100:904-912 (IF 2016 - 2.896)
68. Kolev S, Vassilev D, Kostov K, Todorovska E (2011) Allele variation in loci for adaptive response in Bulgarian wheat cultivars and landraces and its effect on heading date. *Plant Genetic Resources: Characterisation and Utilisation* 9:251-255 (IF 2016 – 0.612)
69. Yediay FE, Andeden EE, Baloch FS, Börner A, Kilian B, Özkan H (2011) The allelic state at the major semi-dwarfing genes in a panel of Turkish bread wheat cultivars and landraces. *Plant Genetic Resources: Characterisation and Utilisation* 9:423-429 (IF 2016 – 0.612)
70. Spanic V, Lemmens M, Drezner G, Dvojkovic K (2011) Interrelations between height of winter wheatgenotypes and resistance to fusarium head blight (FHB) *Romanian Agricultural Research* 28:43-48 (RG IF 2016 – 0.36)
71. Liatukas Ž, Ruzgas V (2011) Coleoptile length and plant height of modern tall and semi-dwarf European winter wheat varieties. *Acta Societatis Botanicorum Poloniae* 80: 197-203 (IF 2016

0.917)

72. Bakshi S, Bhagwat SG (2012) Allelic variations at *Xgwm261* locus, sequence determination and agronomic evaluation in Indian bread wheat genotypes. Cereal Research Communications 40: 34-43 (IF 2016 – 0.496)
73. Asplund L, Leino MW, Hagenblad J (2012) Allelic variation at the *Rht8* locus in a 19th century wheat collection (Review). The Scientific World Journal, v.2012, Article number 385610
74. Gasperini D, Greenland A, Heden R, Harwood W and Griffiths S (2012) Genetic and physiological analysis of *Rht8* in bread wheat: an alternative source of semi-dwarfism with a reduced sensitivity to brassinosteroids. Journal of Experimental Botany 63(12):4419-36 (IF 2016 – 5.83)
75. Maydup ML, Antonietta M, Guiamet JJ, Tambussi EA (2012) The contribution of green parts of the ear to grain filling in old and modern cultivars of bread wheat (*Triticum aestivum* L.): Evidence for genetic gains over the past century. Field Crops Research 134:208-215 (IF 2016 3.048)
76. Petrović S, Marić S, Čupić T, Drezner G, Karsai I (2012) Distribution of allelic variants of hexaploid wheat germplasm at *Xgwm261* and *Ppd-D1* locus. Poljoprivreda 18(2): 25-29
77. Chebotar GA, Chebotar SV, Motsnyy I, Sivolap YuM (2012) The *Rht* and *Ppd-D1* genes in Ukrainian winter bread wheats: effects and distribution. In: European Cereals Genetic Co-operative Newsletter. Proceedings of the 15<sup>th</sup> International EWAC Conference (7-11 November, 2011), pp 62-66
78. Чеботар ГО (2012) Алелі генів короткостебловості *Rht8*, *Rht-B1*, *Rht-D1*, нечутливості до фотoperіоду *Ppd-D1* м'якої пшениці та їх ефекти на агрономічні ознаки. Дисертація на здобуття наукового ступеня канд. біол. наук. – Одеса – 185 с
79. Чеботар ГО, Чеботар СВ, Мощний І, Сиволап ЮМ (2013) Уточнення ступеню зчеплення генів *Rht8* та *Ppd-D1* на 2D хромосомі озимої м'якої пшениці. Цитологія та генетика. том 2013, № 2: 12–17
80. Divashuk MG, Bespalova LA, Vasilyev AV, Fesenko IA, Puzyrnaya OYu, Karlov GI (2013) Reduced height genes and their importance in winter wheat cultivars grown in southern Russia. Euphytica 190: 137-144 (IF 2016 – 1.626)
81. Zhao C, Cui F, Fan Zh, Li J, Ding A, Wang H (2013) Genetic analysis of important loci in the winter wheat backbone parent Aimengniu-V. Australian Journal of Crop Science 7(2):182-188 (RG IF 2016 – 1.02)
82. Chebotar GO, Chebotar SV, Motsnyy II, Sivolap YM (2013) Clarification of the *Rht8-Ppd-D1* gene linkage on the 2D chromosome of winter bread wheat. Cytology and Genetics 47: 70-74 (IF 2016 – 0.324)
83. Zhang J, Dell B, Biddulph B, Drake-Brockman F, Walker E, Khan N, Wong D, Hayden M, Appels R (2013) Wild-type alleles of *Rht-B1* and *Rht-D1* as independent determinants of thousand-grain weight and kernel number per spike in wheat. Molecular Breeding 32( 4):771-783 (IF 2016 – 2.465)
84. Кърцева Т (2014) Проучване ефекта на мутантни гени върху важни биологични качества при домати (*Solanum lycopersicum* L.) и пшеница (*Triticum aestivum* L.). Дисертация за присъждане на образователната и научна степен “Доктор”, ИФРГ-БАН, София ISBN: 978-954-322-722-8
85. Bakshi S, Kumar V, Shitre A, Kishore Das B (2015) Studies to Improve Wheat for High Temperature Stress Areas. American Journal of BioScience 3(3): 64-69
86. Salem KFM (2015) Allelic state at the microsatellite locus *Xgwm261* marking the dwarfing gene *rht8* in Egyptian bread wheat (*Triticum aestivum* L.) genotypes released from 1947 to 2004. Genetika (47): 741-750 (RG IF 2016 – 0.44)
87. Wang Y-Q, Xi W-X, Wang Zh-M, Wang B, Xu X-X, Han M-K, Zhou S-L, Zhang Y-H (2016) Contribution of ear photosynthesis to grain yield under rainfed and irrigation conditions for

winter wheat cultivars released in the past 30 years in North China Plain. Journal of Integrative Agriculture 15: 60345-7 (IF 2016 – 1.042)

**Ganeva G, Landgeva S and Popova Z (2005) Characterization of local accessions of *T. durum* Desf. Bulgaria. Balkan scientific conference – Karnobat, vol1, pp 255-259.**

88. Krasteva L, Stoilova Ts, Varbanova K and Neykov S (2009) Bulgarian Landrace Inventory – Significance and Use. In: Veteläinen M, Negri V and Maxted N (Eds) European landraces on farm conservation, management and use. Biodiversity Technical Bulletin No. 15. Biodiversity International, Rome, Italy, pp 53-68

**Landjeva S, Korzun V, Ganeva G (2006). Evaluation of genetic diversity among Bulgarian winter wheat (*Triticum aestivum* L.) varieties during the period 1925-2003 using microsatellites. Genetic Resources and Crop Evolution 53: 1605-1614**

89. da Costa, CT, Albuquerque ACS, Nascimento Jr A, Marcelino FC, Jorge Fernando Pereira JF (2007) Genetic diversity of Brazilian triticales evaluated with genomic wheat microsatellites. Pesquisa Agropecuaria Brasileira 42 (11): 1577-1586
90. Zhao L, Tian J (2007). AFLP Analysis of genetic diversity of wheat cultivars in Shandong province. Molecular Plant Breeding 5: 403-411
91. Zheleva D, Todorovska E, Christov N, Ivanov P, Ivanova I, Todorov I (2007) Assessing the genetic variation of Bulgarian bread wheat varieties by biochemical and molecular markers. Biotechnology and Biotechnological Equipment 21: 311-321 (IF 2016 – 1.059)
92. Hysing SC (2007) Genetic resources for disease resistance breeding in wheat. Characterization and utilization. PhD thesis, Swedish University of Agricultural Sciences, Alnarp
93. Hysing SC, Säll T, Nybom H, Liljeroth E, Merker A, Orford S, Koebner RMD (2008) Temporal diversity changes among 198 Nordic bread wheat landraces and cultivars detected by retrotransposon-based S-SAP analysis. Plant Genetic Resources: Characterization and Utilization 6(2): 113-125 (IF 2016 – 0.612)
94. Zheleva D (2008) Дисертация за присъждане на образователната и научна степен "Доктор", Агробиоинститут, НЦАН, София
95. Khlestkina EK, Röder MS, Efremova TT, Börner A, Shumny VK (2008) The genetic diversity of old and modern Siberian varieties of common spring wheat as determined by microsatellite markers. Plant Breeding 123: 122 – 127 (IF 2016-1.335)
96. Hu T, Li G, Feng S, Li X, Liu Z, Ru Z, Liu Z (2008) Genetic diversity comparison between wheat cultivars from Chile and Huang-Huai area by SSR Marker. Journal of Henan Agricultural Sciences, 8: 28-31
97. Malik R, Kundu S, Sareen S, Kumar R, Shoran J, Mishra B (2008). KSSR and ISSR markers for assessing DNA polymorphism and genetic diversity among bread wheat varieties of India. In: Appels R, Eastwood R et al. (Eds) Proceedings of 11th International Wheat Genetics Symposium, Brisbane Australia, Sydney Univ Press, vol 2, ISBN:978-1-920899-09-7, pp 421-424
98. Malik R, Chowdhury VK, Sunita J, Kundu S, Jain RK (2008) Applicability of ISSR markers for genetic diversity evaluation in Indian bread wheat genotypes of known origin. Environment and Ecology 26 (1): 126-131
99. Malik, R Chowdhury, VK Jain, S Jain, RK (2008) Microsatellites based genetic diversity analysis of Indian bread wheat genotypes. Environment and Ecology 26 (1): 100-106
100. Khlestkina E, Salina E, Pshenichnikova T, Röder MS, Börner A (2009) Glume coloration in wheat: Allelism test, consensus mapping and its association with specific microsatellite allele. Cereal Research Communications 37:37-43 (IF 2016 – 0.496)
101. Kozlova SA, Khlestkina EK, Salina EA (2009) Specific features in using SNP markers developed for allopolyploid wheat. Russian Journal of Genetics 45: 81-84 (IF 2016 – 0.55)
102. Carvalho A, Lima-Brito J, Maçãs B, Guedes-Pinto H (2009) Genetic diversity and variation among

- botanical varieties of old Portuguese wheat cultivars revealed by ISSR assays. Biochemical Genetics 47:276-294 (IF 2016 - 1.543)
- 103.Horvath A, Didier A, Koenig J, Exbrayat F, Charmet G and Balfourier F (2009) Analysis of diversity and linkage disequilibrium along chromosome 3B of bread wheat (*Triticum aestivum* L.) Theoretical and Applied Genetics 119: 1523-1537 (IF 2016 – 4.132)
- 104.Iqbal N, Tabasum A, Sayed H, Hameed A (2009). Evaluation of genetic diversity among bread wheat varieties and landraces of Pakistan by SSR markers. Cereal Research Communications 37: 489-498 (IF 2016 – 0.496)
- 105.Grądzielewska A, Gruszecka D, Leśniowska-Nowak J (2009) RAPD method in examinations of hybrids of tritcale and *Aegilops juvenalis* (Thell.) Eig. Acta Agrophysica 14(3): 591- 602
- 106.Cheng X, Chen P, Xu H, Zhan K, Dong Zh, Wang X, Cui D (2009) Genetic diversity of wheat germplasm of different countries. Journal of Triticeae Crops 29:803-808
- 107.Cheng X, Li H, Xu H, Dong Zh, Song P, Cui D (2009) Evaluation of Different Sampling Methods on the Genetic Diversity of Wheat Germplasm. Chinese Agricultural Science Bulletin 25(18):388-393
- 108.Mattar MZ (2009) The use of simple sequence repeats for detecting genetic diversity in Egyptian and exotic bread wheat. Egyptian Journal of Genetics and Cytology 38: 325-334
- 109.van de Wouw M, van Hintum T, Kik C, van Treuren R, Visser B (2010) Genetic diversity trends in twentieth century crop cultivars: a meta analysis. Theoretical and Applied Genetics 120:1241–1252 (IF 2016 – 4.132)
- 110.Achtar S, Moualla MY, Kalhout A, Röder MS, MirAli N (2010) Assessment of genetic diversity among Syrian durum (*Triticum* ssp. *durum*) and bread wheat (*Triticum aestivum* L.) using SSR markers. Russian Journal of Genetics 46 (11):1320-1326 (IF 2016 – 0.55)
- 111.Sefera T, Abebie B, Gaur PM, Assefa K, Varshney RK (2011) Characterisation and genetic diversity analysis of selected chickpea cultivars of nine countries using simple sequence repeat (SSR) markers. Crop and Pasture Science 62 (2):177-187 (IF 2016 – 1.804)
- 112.Zhang C, Dong C, He X, Zhang L, Xia X, He Zh (2011) Allelic variants at the TaZds-D1 locus on wheat chromosome 2DL and their association with yellow pigment content. Crop Science 51(4):1580-1590 (IF 2016 – 1.787)
- 113.Petrović S (2011) Genetic diversity of winter bread wheat (*Triticum aestivum* L. ssp *vulgare*). PhD Thesis, Josip Juraj Strossmayer University of Osijek, Croatia
- 114.Guo X, Gao A, Liu W, Yang X, Li X, Li L (2011) Evaluation of genetic diversity, population structure, and linkage disequilibrium among elite Chinese wheat (*Triticum aestivum* L.) cultivars. Australian Journal of Crop Science 5:1167-1172 (RG IF 2016 – 1.02)
- 115.Leonova IN, Budashkina EB, Kalinina NP, Röder MS, Börner A, Salina EA (2011) *Triticum aestivum* – *Triticum timopheevii* introgression lines as a source of pathogen resistance genes. Czech Journal of Genetics and Plant Breeding, 47, (Special Issue): 49–55 (IF 2016 – 0.532)
- 116.Ahmad M (2011) Evaluation of wheat genotypes for salt tolerance based on conventional and molecular approaches, PhD Thesis, Arid Agriculture University Rawalpindi, India
- 117.Menzo VM (2011) Molecular markers validation to develop a Marker Assisted Selection programme in durum wheat (*Triticum turgidum* L. subsp. *durum* (Desf) Husn.). PhD Thesis, Università degli studi della Tuscia – Viterbo
- 118.Karsai I, Vida Gy, Petrovics S, Petcu E, Kobiljski B, Ivanovska S, Bedö Z, Veisz O (2012) Assessment of the spatial genotypic and phenotypic diversity present in the various winter wheat breeding programs in Southeast Europe. Euphytica 186: 139-151 (IF 2016 – 1.626)
- 119.Cheng X, Xu H, Dong Z, Chen F, Zhan K and Cui D (2012) Genetic evolution and utilization of wheat germplasm resources in Huanghuai winter wheat region of China. Pakistan Journal of Botany 44(1): 281-288 (IF 2016 – 0.658)
- 120.Rauf S, Tariq SA, Hassan S (2012) Estimation of pedigree based diversity in Pakistani wheat

(*Triticum aestivum* L.) germplasm. Communications in Biometry and Crop Science 7(1): 14-22

- 121.Španić V, Buerstmayr H, Drezner G (2012) Assessment of genetic diversity of wheat genotypes using microsatellite markers. Periodicum Biologorum 114(1):37-42
- 122.Li H, Zhang H, Liu B, Liu D, Zhang B (2012) Assessing genetic diversity and its changes of bread wheat in Qinghai Province, China, using agronomic traits and microsatellite markers. Biological Agriculture & Horticulture 28: 120-128
- 123.Santos-Garcia MO, de Toledo-Silva G, Sasaki RR, Ferreira TH, Resende RMS, Chiari L, Karia CT, Carvalho MA, Faleiro FG, Zucchi MI, de Souz AP. (2012) Using genetic diversity information to establish core collections of *Stylosanthes capitata* and *Stylosanthes macrocephala*. Genetics and Molecular Biology 35:847-861 (IF 2015 – 1.341)
- 124.Petrović S, Marić S, Čupić T, Drezner G, Karsai I (2012) Assessment of genetic diversity in Croatian winter wheat varieties using SSR and AFLP markers. Poljoprivreda 18(2): 18-24
- 125.Чеботар ГО (2012) Алелі генів короткостебловості Rht8, Rht-B1, Rht-D1, нечутливості до фотoperіоду Ppd-D1 м'якої пшениці та їх ефекти на агрономічні ознаки. Дисертація на здобуття наукового ступеня канд. біол. наук. – Одеса– 185 с
- 126.Sardouie-Nasab S, Mohammadi-Nejad Gh, Nakhoda B (2013). Assessing genetic diversity of promising wheat (*Triticum aestivum* L.) lines using microsatellite markers linked with salinity tolerance. Journal of Plant Molecular Breeding 1(2): 28-39
- 127.Kolesnyk OO, Chebotar SV, Syvolap Yu M, Tsevma VM, Khokhlov OM, Litvinenko MA (2013). The courses of nurseries control and variety investigation systems founded on molecular markers. Збірник наукових праць СГІ–НЦНС. 2013. Вип. 22 (62): 89-99
- 128.Wang Y, Wang Ch, Zhang H, Yue Zh, Liu X, Ji W (2013) Genetic analysis of wheat (*Triticum aestivum* L.) and related species with SSR markers. Genetic Resources and Crop Evolution 60: 1105-1117 (IF 2016 – 1.294)
- 129.Haile JK, Hammer K, Badebo A, Nachit MM, Röder MS (2013) Genetic diversity assessment of Ethiopian tetraploid wheat landraces and improved durum wheat varieties using microsatellites and markers linked with stem rust resistance. Genetic Resources and Crop Evolution 60:513-527 (IF 2016 – 1.294)
- 130.Senturk Akfirat, F, Uncuoglu AA (2013) Genetic diversity of winter wheat (*Triticum aestivum* L.) revealed by SSR markers. Biochemical Genetics 51: 223-229 (IF 2016 – 1.543)
- 131.Drikvand R, Bihamta MR, Najafian G, Ebrahimi A (2013) Investigation of genetic diversity among bread wheat cultivars (*Triticum aestivum* L.) using SSR markers. Journal of Agricultural Science 5(1): 122-129 (IF 2016 - 1.291)
- 132.del Pozo A, Matus I, Serret MD, Araus JL (2014) Agronomic and physiological traits associated with breeding advances of wheat under high-productive Mediterranean conditions. The case of Chile. Environmental and Experimental Botany, 103: 180–189 (IF 2016 - 4.369)
- 133.Dvořáček V, Hermuth J, Dotlačil L, Prohasková A, Hauptvogel P (2014) Changing parameters of Czechoslovak obsolete and modern bread wheat cultivars (*Triticum aestivum* L.) over 90 years. Genetic Resources and Crop Evolution 61, 6, 1159-1171 (IF 2016 – 1.294)
- 134.Khan MK, Pandey A, Choudhary S, Hakki EE, Akkaya MS, Thomas G (2014) From RFLP to DArT: molecular tools for wheat (*Triticum* spp.) diversity analysis. Genetic Resources and Crop Evolution 61, 5, 1001-1032 (IF 2016 – 1.294)
- 135.Arora A, Kundu S, Dilbaghi N, Sharma I, Tiwari R (2014) Population structure and genetic diversity among Indian wheat varieties using microsatellite (SSR) markers. Australian Journal of Crop Science 8(9):1281-1289 (RG IF 2016 – 1.02)
- 136.Kondić-Špika A, Ničić M, Brbaklić L, Trkulja D, Miladinović D, Jocić S, Brdar M, Hristov N (2014) Microsatellites in the analysis of wheat genetic diversity. Genetika 46: 1047-1063 (RG IF 2016 – 0.44)

137. Mollaheydari Bafghi R, Baghizadeh A, Mohammadi-Nejad Gh, Nakhoda B (2014) Assessment of genetic diversity in Iranian wheat (*Triticum aestivum L.*) cultivars and lines using microsatellite markers. *Journal of Plant Molecular Breeding* 2(1): 74-89
138. Salem KFM, Röder MS, Börner A (2015) Assessing genetic diversity of Egyptian hexaploid wheat (*Triticum aestivum L.*) using microsatellite markers. *Genetic Resources and Crop Evolution* 62: 377-385 (IF 2016 – 1.294)
139. Kolesnyk OO, Chebotar SV, Khokhlov OM (2015) Microsatellite markers associated with grain colour of Ukrainian bread wheat varieties. Наукові доповіді Національного університету біоресурсів і природокористування України, № 2. [http://nbuv.gov.ua/j-pdf/Nd\\_2015\\_2\\_10.pdf](http://nbuv.gov.ua/j-pdf/Nd_2015_2_10.pdf).
140. Леонова ИН (2015) Генетический контроль устойчивости к грибным болезням у мягкой пшеницы с интроверсиями от *Triticum timopheevii* Zhuk. Дисертація, Інститут по цитологія и генетика, СО РАН, Новосибирск
141. Novoselovic D, Bentley AR, Šimek R, Dvojkovic K, Sorrells ME, Gosman N, Horsnell R, Drezner G, Šatovic Z (2016) Characterizing Croatian Wheat Germplasm Diversity and Structure in a European Context by DArT Markers. *Frontiers in Plant Science* 7:184. doi: 10.3389/fpls.2016.00184 (IF 2016 - 4.298)
142. Henkrar F, El-Haddoury J, Ouabbou H, Nsarellah N, Iraqi D, Bendaou N, Udupa SM (2016) Genetic diversity reduction in improved durum wheat cultivars of Morocco as revealed by microsatellite markers. *Scientia Agricola* 73(2): 134-141 (RG IF 2016 – 1.15)
143. Orlovskaia OA, Leonova IN, Salina EA, Khotyleva LV (2016) Features of chromosome behavior in meiosis in common wheat lines containing genetic material of tetraploid wheat species. *Russian Journal of Genetics: Applied Research* 6(2): 225-232
144. Soriano JM, Villegas D, Aranzana MJ, García del Moral LF, Royo C (2016) Genetic structure of modern durum wheat cultivars and mediterranean landraces matches with their agronomic performance. *PLoS ONE* 11(8): e0160983. doi:10.1371/journal.pone.0160983 (IF 2016 – 2.806)
145. Kolesnyk OO, Khokhlov OM, Chebotar SV (2016) Associations of alleles of microsatellite markers with agronomical traits of modern bread winter wheat varieties in Southern Ukraine. *Plant Varieties Studying and protection* 3: 19-29
146. Geyer M, Bund A, Albrecht T, Hartl L, Mohler V (2016) Distribution of the fertility-restoring gene *Rf3* in common and spelt wheat determined by an informative SNP marker. *Molecular Breeding* 36: 167. doi:10.1007/s11032-016-0592-6 (IF 2016 – 2.465)
147. Choudhary RC, Sharmja NK, Kumar R, Kumar M (2016) SSR-based genetic diversity assessment among hexaploid wheat (*Triticum aestivum L.*) cultivars. *Indian Journal of Plant Genetic Resources* 29(2): 137-143
148. Meng, L., Xiang, C., Liu, H. et al. (2017) The impact of modern plant breeding on dominant Chinese wheat cultivars (*Triticum aestivum L.*) revealed by SSR and functional markers. *Genetic Resources and Crop Evolution* doi:10.1007/s10722-017-0508-2 (IF 2016 – 1.294)
149. Pražák P, Paczos-Grzeda E (2017) Genetic similarity among *Aegilops kotschy* Boiss. with *Triticum aestivum L.* hybrid lines. *Romanian Agricultural Research*, 34: 15-23 (RG IF 2016 – 0.36)
150. Petrović S, Marić S, Čupić T, Rebekić A, Rukavina I (2017) Assessment of molecular and phenotypic diversity among winter wheat cultivars. *Genetika*, 49: 583-598 (RG IF 2016 – 0.44)
151. Ramya KT, Neelu Jain, Nikita Gandhi, Ajay Arora, Singh PK, Singh Anju M, Singh Gyanendra P, Prabhu KV (2017). Assessing heat stress tolerance and genetic diversity among exotic and Indian wheat genotypes using simple sequence repeats and agro-physiological traits. *Plant Genetic Resources* 15: 208-220 (IF 2016 – 0.612)
- Landjeva S, V Korzun and G Ganeva (2006) Temporal trends in the microsatellite-based genetic diversity of 91 bread wheat varieties released in Bulgaria since 1925. Genetics and Breeding 35 (1-2): 3-10.**
152. Menzo VM (2011) Molecular markers validation to develop a Marker Assisted Selection

programme in durum wheat (*Triticum turgidum* L. subsp. *durum* (Desf) Husn.). PhD Thesis, Università degli studi della Tuscia – Viterbo

**Landjeva S, Korzun V, Tsanev V, Vladova R, Ganeva G (2006). Distribution of the wheat-rye translocation 1RS.1BL among bread wheat varieties of Bulgaria. Plant Breed 125: 102-104**

- 153.Rudnóy Szabolcs R. (2007) Sejtmagi és kloroplasztisz riboszomális DNS szakaszok vizsgálata a hexaploid búzában érsekonsági körében. PhD Thesis, Hungarian Academy of Sciences, Budapest
- 154.Dounia H, Khalfallah N (2008) Comparative analysis of D and R genomes in two lignes (X-*Triticosecale* wittmack) and their genitors (*Secale cereale* L., *Triticum aestivum* L.) by N-banding. Caryologia 61 (3): 245-252 (IF 2016 – 0.516)
- 155.Schlegel R (2008). Current list of wheats with rye introgression. V05-08, 1-14, [http://www.desicca.de/Wheat-rye introgression](http://www.desicca.de/Wheat-rye%20introgression)
- 156.Tang ZX, Fu SL, Ren ZL, Zhang HQ, Yang ZJ, Yan BJ, Zhang HY (2008). Production of a new wheat cultivar with a different 1B.1R translocation with resistance to powdery mildew and stripe rust. Cereal Research Communications 36: 451-460 (IF 2016 – 0.496)
- 157.Singh R, TiwariR, Priyamvada, Gupta RK, Shoran J, MishraB (2009) 1RS.1BL translocation and grain yield as well as bread loaf volume in Indian wheats. Cereal Research Communications 37: 441-448 (IF 2016 – 0.496)
- 158.Mirzaghadri G, Karimzadeh G, Shahsavand H, Jalali M, Baghizadeh A (2009) Wheat-Rye Translocation 1RS.1BL in some bread wheat cultivars of Iran. In: Proc. The 6<sup>th</sup> Nat Biotechol Congress of Iran, 13-15 August 2009, Tehran-Iran, [https://www.researchgate.net/publication/251567248\\_Wheat-Rye\\_Translocation\\_1RS.1BL\\_insome\\_bread\\_wheat\\_cultivars\\_of\\_Iran](https://www.researchgate.net/publication/251567248_Wheat-Rye_Translocation_1RS.1BL_insome_bread_wheat_cultivars_of_Iran)
- 159.Yediay FE, Baloch FS, Kilian B and Özkan H (2010). Testing of rye-specific markers located on 1RS chromosome and distribution of 1AL.RS and 1BL.RS translocations in Turkish wheat (*Triticum aestivum* L., *T.durum* Desf.) varieties and landraces. Genetic Resources and Crop Evolution 57 (1): 119-129 (IF 2016 – 1.294)
- 160.Белан ИА, Рассеева ЛП, Трубачеева НВ, Осадчая ТС, Дорогина ОВ, Жмудь ЕВ, Колмаков ЮВ, Блохина НП, Кравцова ЛА, Першина ЛА (2010). Особенности хозяйственно ценных признаков линий сорта яровой мягкой пшеницы Омская 37, несущих пшенично-ржаную транслокацию 1RS.1BL. Вестник ВОГиС, Том 14, № 4: 632-640
- 161.Trubacheeva NV, Rosseeva LP, Belan IA, Osadchaya TS, Kravtsova LA, Kolmakov YV, Blokhina NP, Pershina LA (2011). Characteristics of common wheat cultivars of West Siberia carrying the wheat-rye 1RS.1BL translocation. Russian Journal of Genetics 47: 13-18 (IF 2016 – 0.55)
- 162.Tang ZX, Ross K, Ren ZL, Yang ZJ, Zhang HY, Chikmawati T, Miftahudin and Gustafson JP (2011) Secale. In: Chittaranjan Kole (Ed) Wild Crop Relatives: Genomic and Breeding Resources. Cereals. Springer
- 163.Mirzaghadri G, Zeinali G, Rafiepour M, Karimzadeh G (2011) Wheat-rye translocation in Iranian bread wheat cultivars and their ion distribution in response to salinity stress. Journal of Agricultural Science and Technology 13, Issue Suppl., 1163-1172 (IF 2016 – 0.813)
- 164.Belan IA, Rosseeva LP, Trubacheeva NV, Osadchaya TS, Dorogina OV, Zhmud' EV, Kolmakov YuV, Blokhina NP, Kravtsova LA and Pershina LA (2011) Important agronomic traits of Omskaya 37 lines of spring wheat cultivar carrying wheat-rye translocation 1RS.1BL. Russian Journal of Genetics: Applied Research 1: 241-247
- 165.Zhao C, Cui F, Wang X, Shan S, Li X, Bao Y, Wang H (2012) Effects of 1BL/1RS translocation in wheat on agronomic performance and quality characteristics. Field Crops Research 127: 79-84 (IF 2016 3.048)
- 166.(Identification of Chromosome Segments of Rye 1RS in Aegilops Genus using DNA Specific Molecular Marker) (2012) – иранско списание, с ирански юероглифи [https://journals.ut.ac.ir/article\\_29046\\_43.html](https://journals.ut.ac.ir/article_29046_43.html)

- 167.Tabibzadeh N, Karimzadeh G, Naghavi M R (2013) Distribution of 1AL.1RS and 1BL.1RS wheat-rye translocations in Iranian wheat, using PCR based markers and SDS-PAGE. Cereal Research Communications 41(3): 458-467 (IF 2016 – 0.496)
- 168.Zhou J, Han C, Cao H, Zhen S , Yu Z, Li X, Ma W, Yan Y (2013) Fast identification of wheat 1BL.1RS translocation by reversed-phase ultra-performance liquid chromatography (RP-UPLC). Crop and Pasture Science 64(9): 865-873 (IF 2016 – 1.804)
- 169.Bagherikia S, Karimzadeh G, Naghavi M R (2013) Identification of Rust-Resistance Genes (Lr26, Sr31, Yr9) Using Specific PCR. Crop Biotechnology Autumn and Winter 5: 129-138
- 170.Tahir R, Bux H, Kazi AG, Rasheed A, Napar AA, Ajmal SU, Mujeeb-Kazi A (2014) Evaluation of Pakistani Elite Wheat Germplasm for T1BL.1RS Chromosome Translocation. Journal of Agricultural Science and Technology 16:421-432 (IF 2016 – 0.813)
- 171.Bagherikia S, Karimzadeh G, Naghavi M R (2014) Distribution of 1AL.1RS and 1BL.1RS wheat-rye translocations in *Triticum aestivum* using specific PCR. Biochemical Systematics and Ecology 55: 20-26 (IF 2016 -0.929)
- 172.Hammami, R, Jouve N, Soler C, Frieiro E, González JM (2014) Genetic diversity of SSR and ISSR markers in wild populations of *Brachypodium distachyon* and its close relatives *B. stacei* and *B. hybridum* (Poaceae). Plant Systematics and Evolution 300: 229 (IF 2016 - 1.239)
- 173.Motsny II, Sudarchuk LV, Galaev AV, Chebotar SV (2014) Application of PCR markers for detecting 1BL.1RS wheat-rye chromosome translocations and (1B)1R substitutions. Factors in experimental evolution of organisms 15: 264-268
- 174.Силкова ОГ, Логинова ДБ, Иванова (Кабаненко) ЮН, Бондаревич ЕБ, Соловей ЛА, Штык ТИ, Дубовец НИ (2014) Интрогрессия хроматина ржи в геном мягкой пшеницы: цитогенетические аспекты. Вавиловский журнал генетики и селекции, 2014, Том 18, № 4/1: 630-642
- 175.Hakki EE, Dogrur N, Pandey A, Khan MK, Hamurcu M, Kayis SA, GEzgin S, Ölmez F, Akkaya MS (2014) Molecular and elemental characterization of selected turkish durum wheat varieties, Not Bot Horti Agrobo, 42, 2:431-439 (IF 2016 – 0.480)
- 176.Guberac S, Petrović S, Guberac V, Marić S (2015) 1RS translocation in Croatian winter wheat varieties. Poljoprivreda/Agriculture, 21(1): 8-14
- 177.Uzma, Kubra Gh, Gul A, Mujeeb-Kazi A (2015). Use of alien diversity to combat some major biotic stresses in *Triticum aestivum* L. In: Hakeem K R (Ed) Crop Production and Global Environmental Issues, Springer, pp. 319-348
- 178.Naseeb Ullah, Niaz Ali, Muhammad Iqbal, Aziz-ud-Din, Azhar Hussain Shah, Inayat Ur Rahman, Habib Ahmad, Inamullah, Ghulam Muhammad Ali (2016) Markers assisted selection for multiple stripe rust resistance genes in spring bread wheat lines. International Journal of Biosciences 8 (3): 63-74
- 179.Моргун БВ, Лялько ІІ, Степаненко АІ, Бавол АВ, Великожон ЛГ (2016) Цитологічна та молекулярно-генетична характеристика сортів озимої м'якої пшениці, отриманих за інтрогресивної селекції та м. Фактори експериментальної еволюції організмів, Том 19, 167-171
- 180.Lan C, Hale IL, Herrera-Foessel SA, Basnet BR, Randhawa MS, Huerta-Espino J, Dubcovsky J, Singh R (2017) Characterization and mapping of leaf rust and stripe rust resistance loci in hexaploid wheat lines UC1110 and PI610750 under Mexican environments. Frontiers in Plant Science 8:1450 (IF 2016 - 4.298)
- 181.Golkari S (2017) The evaluation of rye 1RS translocation distribution in some Iranian wheat promising genotypes and wheat cultivars using molecular markers. Modern Genetics, 12: 133-141
- 182.Duan Q, Wang YY, Qiu L, Ren TH, Li Z, Fu SL, Tang ZX (2017) Physical location of new PCR-based markers and powdery mildew resistance gene(s) on rye (*Secale cereale* L.) chromosome 4 using

4R dissection lines. Frontiers in Plant Science, <https://doi.org/10.3389/fpls.2017.01716> (IF 2016 - 4.298)

**Landjeva S, Ganeva G (2006). Changes in the seedling growth parameters in three common wheat (*Triticum aestivum* L) cultivars subjected to drought stress and subsequent re-hydration. Field Crop Studies 3: 185-190**

- 183.Kosturkova G, Todorova R, Sakthivelu G, Akitha Devi MK, Giridhar P, Rajasekaran T, Ravishankar GA (2008) Response of Bulgarian and Indian soybean genotypes to drought and water deficiency in field and laboratory conditions. General and Applied Plant Physiology 34:239-250
- 184.Radaei Alamoli Z, Akbari Gh, Abdollahian-Noghabi M (2013) Effect of drought stress in the field and the relationship between an in vitro method (Polyethylene glycol 6000) for screening sugar beet genetic resources. International Journal of AgriScience 3(11): 838-850

**Landjeva S, Korzun V, Börner A (2007). Molecular markers: actual and potential contributions to wheat genome characterization and breeding. Euphytica 156: 271-296**

- 185.Araus JL, Slafe, GA, Royo C, Serret MD (2008) Breeding for yield potential and stress adaptation in cereals. Critical Reviews in Plant Science 27:377–412 (IF 2016 - 6.825)
- 186.Gupta PK, Mir RR, Mohan A, Kumar J (2008) Wheat genomics: present status and future prospects. In: Gupta PK and Yunbi Xu (Eds) Genomics of Major Crops and Model Plant Species, 36 pages. Article ID 896451; DOI:10.1155/2008/896451
- 187.Halloran GM, Ogbonnaya FC, Lagudah ES (2008) *Triticum (Aegilops) tauschii* in the natural and artificial synthesis of hexaploid wheat. Australian Journal of Agricultural Research 59 (5): 475-490 (IF 2016 – 1.804)
- 188.Hao Y, Liu A, Wang Y, Feng D, Gao J, Li X, Liu S, Wang H (2008) Pm23: A new allele of *Pm4* located on chromosome 2AL in wheat. Theoretical and Applied Genetics 117: 1205-1212 (IF 2016 – 4.132)
- 189.Pánková K, Milec Z, Leverington-Waite M, Chebotar S, Snape JW (2008) Characterization of inter-varietal chromosome substitution lines of wheat using molecular markers. Czech Journal of Genetics and Plant Breeding 44: 22–29 (IF 2016 – 0.532)
- 190.Navakode S (2008) Molecular mapping of quantitative trait loci (QTL) controlling aluminium tolerance in wheat and barley. PhD Thesis, Martin-Luther-Universität Halle-Wittenberg
- 191.Li G-R, Liu C, Zeng Z-X, Jia J-Q, Zhang T, Zhou J-P, Ren Z-L, Yang Z-J (2009) Identification of  $\alpha$ -gliadin genes in *Dasypyrum* in relation to evolution and breeding. Euphytica 165: 155-163 (IF 2016 – 1.626)
- 192.Brdar-Jokanović M (2009) Phenotypic and molecular variability of boron tolerance in wheat, PhD Thesis, University of Novi Sad
- 193.Francki MG, Walker E, Crawford AC, Broughton S, Ohm HW, Barclayl, Wilson RE and McLean R (2009) Comparison of genetic and cytogenetic maps of hexaploid wheat (*Triticum aestivum* L.) using SSR and DArT markers. Molecular Genetics and Genomics 281:181-191 (IF 2016 - 2.979)
- 194.Hanzalová A, Sumíková T and Bartoš P (2009) Determination of leaf rust resistance genes Lr10, Lr26 and Lr37 by molecular markers in wheat cultivars registered in the Czech Republic. Czech Journal of Genetics and Plant Breeding, 45 (2): 79–84 (IF 2016 – 0.532)
- 195.Petrović S, Marić S, Guberac V, Drezner G, Eđed A (2009) Varijabilnost kvantitativnih svojstava germplazme ozime pšenice. Proc. 44th Croatian & 4th International Symposium on Agriculture, February 16 - 20, 2009, Opatija, Croatia, pp 365-369
- 196.Prasad B, Babar MA, Xu XY, Bai, GH, Klatt AR (2009) Genetic diversity in the U.S. hard red winter wheat cultivars as revealed by microsatellite markers. Crop and Pasture Science 60:16-24 (IF 2016 – 1.804)
- 197.Quincke MC (2009) Phenotypic Response and Quantitative Trait Loci for Resistance to *Cephalosporium gramineum* in Winter Wheat, PhD Thesis, Oregon State University
- 198.Wanjugi H, Coleman-Derr D, Huo N, Kianian SF, Luo MC, Wu J, Anderson O, Gu YQ (2009) Rapid

- development of PCR-based genome specific repetitive DNA junction markers in wheat. *Genome* 52: 576–587
199. Hammer K, Diederichsen A (2009) Evolution, Status and Perspectives for Landraces in Europe. In: Veteläinen M, Negri V and Maxted N (Eds.). European landraces on farm conservation, management and use. Bioversity Technical Bulletin No. 15. Biodiversity International, Rome, Italy, pp 23-44
200. Kobiljski B, Denčić S and Kondić-Špika A (2009). Integration of conventional and molecular wheat breeding strategies. In: Proc of International Conference 'Conventional And Molecular Breeding Of Field And Vegetable Crops' Novi Sad, 24-27 November 2008, pp 587-589
201. Holzapfel JJ (2009) Kartierung von Resistenzloci gegen Ährenfusarium im europäischen Winterweizen unter Berücksichtigung des Kurzstrohgens Rht-D1. PhD Thesis, TU-München
202. Xue F, Zhai W-W, Duan X-Y, Zhou Y-L and Wan-Quan JI (2009) Microsatellite mapping of powdery mildew resistance gene in wheat landrace Xiaobaidong. *Acta Agronomica Sinica* 35(10): 1806–1811 (RG IF 2016 – 0.12)
203. Hovhannisan NA, Yesayan AH (2010) Genetic diversity of *Aegilops tauschii* populations in Armenia. *Biological Journal of Armenia* 62 (4): 59-65
204. Torres AM (2010) Application of Molecular Markers for Breeding Disease Resistant Varieties in Crop Plants. In: S Mohan Jain and DS Brar (Eds) Molecular Techniques in Crop Improvement, 2nded, Springer, Part 2, Chapter 8, pp 185-205 (DOI 10.1007/978-90-481-2967-6\_8)
205. Paux E, Faure S, Choulet F, Roger D, Gauthier V, Martinant J-P, Sourdille P, Balfourier F, Le Paslier M-C, Chauveau A, Cakir M, Gandon B, Feuillet C (2010) Insertion site-based polymorphism markers open new perspectives for genome saturation and marker-assisted selection in wheat. *Plant Biotechnology Journal* 8 (2): 196-210 (IF 2016 – 7.443)
206. Gupta PK, Langridge P, Mir RR (2010) Marker-assisted wheat breeding: present status and future possibilities. *Molecular Breeding* 26(2): 145-161 (IF 2016 – 2.465)
207. Niu J-S, Jia H-Y, Yin J, Wang B-Q, Ma Z-Q, Shen T-M (2010) Development of an STS marker linked to powdery mildew resistance genes *PmLK906* and *Pm4a* by gene chip hybridization. *Agricultural Sciences in China* 9 (3): 331-336
208. Bhullar NK, Mackay M, Keller B (2010) Genetic diversity of the *Pm3* powdery mildew resistance alleles in wheat gene bank accessions as assessed by molecular markers. *Diversity* 2: 768-786
209. Sumíková T and Hanzalová A (2010). Multiplex PCR assay to detect rust resistance genes *Lr26* and *Lr37* in wheat. *Czech Journal of Genetics and Plant Breeding*, 46 (2): 85–89 (IF 2016 – 0.532)
210. Schubert A (2010). Quantitative trait loci analysis in spring wheat comparing two advanced backcross populations derived from an exotic wheat accession. PhD Thesis, Rheinischen Friedrich-Wilhelms-University, Bonn, Germany
211. Bibi S, Dahot MU, Nizamani GS, Khan IA, Khatri A, Naqvi MH, Oad FC, Burio UA (2010) Molecular marker assisted selection for drought tolerant wheat genotypes. *Pakistan Journal of Botany* 42(4): 2443-2452 (IF 2016 – 0.658)
212. Мельникова НВ (2010) Мировое разнообразие твердой пшеницы (*Triticum durum* Desf.) по аллелям глиадинкодирующих локусов. Диссертация кбн, Москва
213. Risser P (2010) Mapping of quantitative trait loci (QTL) for adult-plant resistance to *Septoria tritici* in five wheat populations (*Triticum aestivum* L.). PhD Thesis Universität Hohenheim, Germany
214. Gupta PK, Kumar J, Mir RR, Kumar A (2010). Marker-assisted Selection as a Component of Conventional Plant Breeding. In: J Janick (Ed) *Plant Breeding Reviews* volume 33, John Wiley & Sons, Inc. Hoboken, New Jersey, pp 145-218
215. Andrade F, Abbate P, Otegui M, Cirilo A, Cerrudo A (2010) Ecophysiological basis for crop management. *The Americas Journal of Plant Science and Biotechnology* 4: 23-34

- 216.Arief VN, Kroonenberg PM, DeLacy IH, Dieters MJ, Crossa J, Dreisigacker S, Braun H-J, Basford KE (2010) Construction and three-way ordination of the wheat genome atlas. *Journal of Applied Probability and Statistics* 5 (2): 95-118.
- 217.Reynolds M, Bonnett D, Chapman SC, Furbank RT, Manès Y, Mather DE and Parry MAJ (2011) Raising yield potential of wheat. I. Overview of a consortium approach and breeding strategies. *Journal of Experimental Botany* 62: 439-452 (IF 2016 – 5.83)
- 218.Yu F, Wang B-H, Feng S-P, Wang J-Y, Li W-G, Wu Y-T (2011). Development, characterization, and cross-species/genera transferability of SSR markers for rubber tree (*Hevea brasiliensis*). *Plant Cell Report* 30(3):335-344 (IF 2016 - 2.869)
- 219.Lata C, Prasad M (2011). Role of DREBs in regulation of abiotic stress responses in plants. *Journal of Experimental Botany* 62: 4731-4748 (IF 2016 – 5.83)
- 220.Paux E, Sourdille P, Mackay I, Feuillet C (2011). Sequence-based marker development in wheat: Advances and applications to breeding, *Biotechnology Advances* 30 (5): 1071-1088 (IF 2016 – 10.597)
- 221.Yin D-D, An D-G, Li L-H, Xu H-X (2011). Application of molecular marker techniques in rye research. *Chinese Journal of Eco-Agriculture* 19: 477-483
- 222.Ahmad M (2011) Evaluation of wheat genotypes for salt tolerance based on conventional and molecular approaches, PhD Thesis, Arid Agriculture University Rawalpindi, Pakistan
- 223.Ovesná J, Janská A, Zelenková S, Marík P (2011) Chapter 2. High-Throughput Approaches for Characterization and Efficient Use of Plant Genetic Resources. In: Benkeblia N (Ed) Sustainable Agriculture and New Biotechnologies, CRC Press 2011, pp 23-39
- 224.Ortiz R (2011) Advances in wheat genetic enhancement for global food production. Chapter 4 in: Almeida MT (Ed.) Wheat: Genetics, Crops and Food Production. Nova Science Publishers, Inc.pp 153-178
- 225.Agarwal R, Kharbikar LL, Renu (2011) Identification of Bipolaris sorokiniana- responsive differential transcripts in wheat (*Triticum aestivum* L.). *Indian Phytopathology* 64 (1) : 24-27
- 226.Kavanagh VB (2012) Environmental biosafety of field scale GM triticale (xTriticosecale Wittmack) cultivation for bioindustrial applications. PhD thesis, University of Alberta, Edmonton
- 227.Takashi Nakatsuka, Misa Saito, Yuka Sato-Ushiku, Eri Yamada, Takashi Nakasato, Nobue Hoshi, Kazumichi Fujiwara, Takashi Hikage and Masahiro Nishihara (2012) Development of DNA markers that discriminate between white- and blue-flowers in Japanese gentian plants. *Euphytica* 184 (3): 335-344 (IF 2016 – 1.626)
- 228.Huang BE, George AW, Forrest KL, Kilian A, Hayden AJ, Morell MK, Cavanagh CR (2012) A multiparent advanced generation inter-cross population for genetic analysis in wheat. *Plant Biotechnology Journal* 10: 826-839 (IF 2016 – 7.443)
- 229.Khlestkina EK (2012) Molecular methods for analyzing the structure-function organization of genes and genomes in higher plants. *Russian Journal of Genetics: Applied Research* 2(3): 243-251
- 230.Gao H, Zhu F, Jiang Y, Wu J, Yan W, Zhang Q, Jacobi A, Cai S (2012) Genetic analysis and molecular mapping of a new powdery mildew resistant gene Pm46 in common wheat. *Theoretical and Applied Genetics* 125(5):967-73 (IF 2016 – 4.132)
- 231.Trkulja D, Kondić-Špika A, Brbaklić L, Kobiljski B, Hristov N (2012) Marker-trait associations for spike-related characters in a doubled haploid population of wheat. *Romanian Agricultural Research* 29: 9-16 (RG IF 2016 – 0.36)
- 232.Schneider A, Molnár-Láng M (2012) Detection of various U and M chromosomes in wheat-*Aegilops biuncialis* hybrids and derivatives using fluorescence in situ hybridisation and molecular markers. *Czech Journal of Genetics and Plant Breeding* 48 (4):169–177 (IF 2016 – 0.532)
- 233.Chrpová J, Šíp V, Bartoš P, Hanžalová A, Palicová J, Štočková L, Čejka L, Bížová I, Laml P, Nováček

- T, Horčička P (2012) Results of the Czech National Ring Testsof Disease Resistance in Wheat. Czech Journal of Genetics and Plant Breeding 48 (4): 189–199 (IF 2016 – 0.532)
234. Tocho EF, Marino de Remes Lenicov AM, Castro AM (2012) Assessment of resistance against *Schizaphis graminum* (Hemiptera:Aphididae) in barley. Rev. Soc. Entomol. Argent. 71(1-2): 1-10
235. Gedye KR, Gonzalez-Hernandez JL, Owens V, Boe A (2012) Advances towards a Marker-Assisted Selection Breeding Program in Prairie Cordgrass, a Biomass Crop. International Journal of Plant Genomics, Vol. 2012, Article ID 313545, doi:10.1155/2012/313545
236. Леонова ИН (2013) Молекулярные маркеры: использование в селекции зерновых культур для идентификации, интродукции и пирамидирования генов, Вавиловский журнал генетики и селекции, 17, 2, 314-325
237. Diederichsen A, Solberg S, Jeppson S (2013) Morphological changes in Nordic spring wheat (*Triticum aestivum* L.) landraces and cultivars released from 1892 to 1994. Genetic Resources and Crop Evolution 60: 569-585 (IF 2016 – 1.294)
238. Timonova EM, Leonova IN, Röder MS, Salina EA (2013) Marker-assisted development and characterization of a set of *Triticum aestivum* lines carrying different introgressions from the *T. timopheevii* genome. Molecular Breeding 31(1): 123-136 (IF 2016 – 2.465)
239. Goutam U, Kukreja, Tiwari, Chaudhury A, Gupta RK, Dholakia BB, Yadav R (2013) Biotechnological approaches for grain quality improvement in wheat: Present status and future possibilities. Australian Journal of Crop Science 7: 469-483 (RG IF 2016 – 1.02)
240. Tripathi A, Aggarwal R, Yadav A (2013) Differential expression analysis of defense-related genes responsive to *Tilletia indica* infection in wheat. Turkish Journal of Biology 37(5):606-613 (IF 2016 – 1.038)
241. Lado B, Matus I, Rodriguez A, Inostroza L, Poland J, Belzile F, del Pozo A, Quincke M, von Zitzewitz J (2013) Increased genomic prediction accuracy in wheat breeding through spatial adjustment of field trial data. Genes, Genomes, Genetics; 3(12): 2105-14
242. Токубаева А, Шулембаева К (2013) Идентификация генов устойчивости к бурой ржавчине Lr26, Lr28, Lr34, Lr37c помошью STS и SCAR маркеров у мягкой пшеницы (*Triticum aestivum* L.). KazNU Bulletin. Biology series №1 (57):158-164
243. Du WL, Wang J, Wang LM, Pang YH, Wu J, Zhao JX, Yang QH, Chen XH (2013) A novel SCAR marker for detecting *Psathyrostachys huashanica* Keng chromatin introduced in wheat. Genetics and Molecular Research 12 (4):4797-4806
244. Edwards D, Gupta PK (2013) Sequence Based DNA Markers and Genotyping for Cereal Genomics and Breeding. In: Gupta PK, Varshney R (Eds) Cereal Genomics II, pp 57-76. Springer Netherlands
245. Leonova IN (2013) Molecular markers: Implementation in crop plant breeding for identification, introgression and gene pyramiding. Russian Journal of Genetics: Applied Research 3(6): 464-473
246. Ali N, Iqbal M, Yasmin S, Ahmed I, Asif M (2013) Genetic variation at loci controlling quality traits in spring wheat. Pakistan Journal of Agricultural Science 50(4), 637-647
247. Dura SAM, Duwayri MA, Nachit MM (2013) Detection of molecular markers associated with yield and yield components in durum wheat (*Triticum turgidum* L. var. durum Desf.) under drought conditions. African Journal of Agricultural Research 8(19): 2118-2128
248. Edae EA (2013) Association mapping for yield, yield components and drought tolerance-related traits in spring wheat grown under rainfed and irrigated conditions. PhD Thesis, Colorado State University, USA
249. Dura S, Duwayri M, Nachit M, Al Sheyab F (2014) Detection of molecular markers associated with yield and yield components in durum wheat (*Triticum turgidum* L. var. *durum*) under saline conditions. Crop and Pasture Science 64(10): 957-964 (IF 2016 – 1.804)

- 250.Nielsen NH, Backes G, Stougaard J, Andersen SU, Jahoor A (2014) Genetic Diversity and Population Structure Analysis of European Hexaploid Bread Wheat (*Triticum aestivum* L.) Varieties. PLoS ONE 9(4): e94000. doi:10.1371/journal.pone.0094000 (IF 2016 – 2.806)
- 251.Khan MK, Pandey A, Choudhary S, Hakki EE, Akkaya MS, Thomas G (2014) From RFLP to DArT: molecular tools for wheat (*Triticum* spp.) diversity analysis. Genetic Resources and Crop Evolution 61:1001-1032 (IF 2016 – 1.294)
- 252.Engelmann U (2014) Kartierung und züchterische Nutzung von Resistzenzen gegen die Weizenblattdürre (*Pyrenophora tritici-repentis*). PhD Thesis, Julius Kühn-Institut, Quedlinburg, Germany
- 253.Kondić-Špika A, Ničić M, Brbaklić L, Trkulja D, Miladinović D, Jocić S, Brdar M, Hristov N (2014) Microsatellites in the analysis of wheat genetic diversity. Genetika 46: 1047-1063 (RG IF 2016 – 0.44)
- 254.Леонова ИН, Орловская ОА, Родер МС, Несторов МА, Будашкина ЕБ (2014). Молекулярно-генетическое разнообразие интrogессивных линий мягкой пшеницы (*T. aestivum* /*T. timopheevii*). Вавиловский журнал Генетики и Селекции т. 18, № 4/1: 681-690.
- 255.Huggins, TD (2014). Understanding the Genetic Interactions that Regulate Heat and Drought Tolerance in Relation to Wax Deposition and Yield Stability in Wheat (*Triticum Aestivum* L.). Doctoral dissertation, Texas A & M University
- 256.Kaur P (2015) Physical mapping of chromosome 2AL of hexaploid wheat and generation and mapping of EST based SNPs in *Triticum monococcum*. PhD Thesis, Punjab Agricultural University, India
- 257.Brbaklić L, Trkulja D, Kondić-Špika A, Hristov N, Denčić S, Mikić S, Tomičić M, Kobiljski B (2015) Genetic associations in the detection of QTLs for wheat spike-related traits. Pesquisa Agropecuaria Brasileira 50(2): 149-159
- 258.Abičić I, Lalić A , Šimić G, Šimon S, Pejić I (2015) Genetska raznolikost germplazme ječma Poljoprivrednog instituta Osijek temeljena na molekularnim podatcima. In: Pospišil M (Ed) Proc. 50th Croatian and 10th International Symposium on Agriculture, 16-20 Feb 2015, Opatija, Croatia, pp 211–215
- 259.Aalami A, Mandoulakani BA, Azizi H, Masoumi F, Safiyar S, Karami N (2015) R-RAP: A new marker for genetic characterization and evaluation of relationships among different *Aegilops* species. Crop Breeding Journal 4(1):15-21.
- 260.Lu Y, Wu X, Yao M, Zhang J, Liu W, Yang X, Li X, Du J, Gao A, Li L (2015) Genetic mapping of a putative *Agropyron cristatum*-derived powdery mildew resistance gene by a combination of bulked segregant analysis and single nucleotide polymorphism array. Molecular Breeding 35: DOI: 10.1007/s11032-015-0292-7 (IF 2016 – 2.465)
- 261.Chaabane R, A Saidi , M Rouissi, E Ben Naceur, K Khamassi, C Mejri, M'b Ben Naceur (2015) Phenotyping and genotyping for improving salt tolerance of Tunisian durum wheat cultivars. Options Méditerranéennes,
- 262.Leonova IN, Orlovskaya OA, Röder MS, Nestorov MA, Budashkina EB (2015) Molecular diversity of common wheat introgression lines. (*T. aestivum* /*T. timopheevii*) Russian Journal of Genetics: Applied Research 5 (3):191–197.
- 263.Goutam U, Kukreja S, Yadav R, Salaria N, Thakur K and Goyal AK (2015) Recent trends and perspectives of molecular markers against fungal diseases in wheat. Frontiers in Microbiology 6:861. doi: 10.3389/fmicb.2015.00861 (IF 2016 - 4.165)
- 264.Tian J, Chen J, Chen G, Wu P, Zhang H, Zhao Y (2015) Molecular Markers of Quality Traits and Their Applications. Chapter in: Genetic Analyses of Wheat and Molecular Marker-Assisted Breeding, Volume 2, pp 259-279

- 265.Tian J, Chen J, Chen G, Wu P, Zhang H, Zhao Y (2015) The Concept and Research Progress of MAS. Chapter in: Genetic Analyses of Wheat and Molecular Marker-Assisted Breeding, Volume 2, pp 179-185
- 266.Han B, Wang C, Tang Z, Zhao X (2015) Genome-Wide Analysis of Microsatellite Markers Based on Sequenced Database in Chinese Spring Wheat (*Triticum aestivum* L.) PLOS ONE 10(11):E0141540 (IF 2016 – 2.806)
- 267.Колесник ОО, Чеботар СВ, Хохлов ОМ, Файт ВІ (2015). Порівняння методів диференціації та ідентифікації сортів пшеници м'якої озимої за агрономічними ознаками і морфометричними параметрами зерна та за алельним складом мікросателітних локусів. Вісник Харківського національного аграрного університету серія біологія, вип. 1 (34), 46-54
- 268.Коршунова АД (2015) Влияние генов короткостебельности на хозяйственно- ценные признаки яровой тритикале. Диссертация на соискание ученой степени кандидата биологических наук, ФГБОУ ВО «Российский Государственный Аграрный Университет – МСХА имени К.А. Тимирязева» Москва
- 269.Brbaklić L. (2015) Molecular and association analysis of spike - related traits in wheat (*Triticum aestivum* L.). PhD Thesis, University of Belgrade
- 270.Trkulja D (2015) Association analysis of agronomical traits and SSR loci of wheat. PhD Thesis, University of Belgrade
- 271.Хлесткина ЕК, Пшеничникова ТА, Усенко НИ, Отмахова ЮС (2016) Перспективные возможности использования молекулярно-генетических подходов для управления технологическими свойствами зерна пшеницы в контексте цепочки «зерно – мука – хлеб». Вавиловский журнал генетики и селекции. 20: 511-527.
- 272.Erayman M, İlhan E, Eren AH, Güngör H, Akgöl B (2016) Diversity analysis of genetic, agronomic and quality characteristics of bread wheat (*Triticum aestivum* L.) cultivars grown in Turkey. Turkish Journal of Agriculture and Forestry 40(1): 83-94 (IF 2016 – 1.288)
- 273.Khlestkina, EK, Shumny VK (2016) Prospects for application of breakthrough technologies in breeding: The CRISPR/Cas9 system for plant genome editing. Russian Journal of Genetics 52: 676-687 (IF 2016 – 0.55)
- 274.Štrbanović RT (2016) Identification of alfalfa cultivars application of molecular markers in the initial phases of development plants. PhD Thesis, Univ of Belgrade
- 275.Šušek A (2016) Perspectives of Christmas rose (*Helleborus niger* L.) genetic improvement. Agricultura 13 (1-2): 11-19
- 276.Gerard GS (2017) Identificación de genes de resistencia a mancha de la hoja y otros caracteres de importancia agronómica en trigo. PhD Thesis, Universidad Nacional de La Plata
- 277.Ya N, Raveendar S, Bayarsukh N, Ya M, Jung-Ro Lee, Lee K-J, Shin M-J, Cho G-T, Ma K-H, Lee G-A (2017) Genetic diversity and population structure of mongolian wheat based on SSR markers: implications for conservation and management. Plant Breeding and Biotechnology 5(3):213-220
- 278.Khlestkina EK, Pshenichnikova TA, Usenko NI, Otmakhova YuS (2017) Promising opportunities of using molecular genetic approaches for managing wheat grain technological properties in the context of the “grain–flour–bread” chain. Russian Journal of Genetics: Applied Research, 7: 459-476
- 279.Jiang M, Xaio ZQ, Fu SL, Tang ZX (2017) FISH karyotype of 85 common wheat cultivars/lines displayed by ND-FISH using oligonucleotide probes. Cereal Research Communication <https://doi.org/10.1556/0806.45.2017.049> (IF 2016 – 0.496)

Landjeva S, Neumann K, Lohwasser U, Börner A (2008). Molecular mapping of genomic regions associated with wheat seedling growth under osmotic stress. Biol Plant 52: 259-266

- 280.Kosturkova G, Todorova R, Sakthivelu G, Akitha Devi MK, Giridhar P, Rajasekaran T, Ravishankar GA (2008). Response of Bulgarian and Indian soybean genotypes to drought and water deficiency in field and laboratory conditions. General and Applied Plant Physiology 34:239-250
- 281.Zhuang YL, Ren GJ, Zhu Y, Hou GH, Qu X, Li ZX, Yue GD, Zhang JR (2008). Transcriptional profiles of immature ears and tassels in maize at early stage of water stress. Biologia Plantarum 52: 754-758 (IF 2016 – 1.551)
- 282.Maccaferri M, Sanguineti MC, Giuliani S, Tuberosa R (2009) Genomics of Tolerance to Abiotic Stress in the Triticeae. In: Muehlbauer GJ and Feuillet C (eds) Genetics and Genomics of the Triticeae. Plant Genetics and Genomics: Crops and Models vol 7, Chapter 18, pp 481-558. Springer New York
- 283.Yin Z-T, Song H-N, Meng F-F, Xu X-M and Yu D-Y (2010). QTL mapping for photosynthetic gas-exchange parameters in soybean. Acta Agronomica Sinica 36(1): 92–100 (RG IF 2016 – 0.12)
- 284.Hamayun M, Khan SA, Shinwari ZK, Khan AL, Ahmad N and Lee I-J (2010) Effect of polyethylene glycol induced drought stress on physio-hormonal attributes of soybean. Pakistan Journal of Botany 42: 977-986 (RG IF 2016 – 0.94)
- 285.Mirzaghaderei G, Karimzadeh G, Hassani HS, Jalali-Javaran M and Baghizadeh A (2010) Cytogenetic analysis of hybrids derived from wheat and Tritipyrum using conventional staining and genomic in situ hybridization. Biologia Plantarum 54 (2): 252-258 (IF 2016 – 1.551)
- 286.Li Zh, Yuan Q, Shi C, Chen J, Han S, Tian J (2010) QTL mapping for coleoptile length and root length in wheat seedling. Molecular Plant Breeding 8 (3):460-468
- 287.Khalil Zaynali Nezhad (2010). Genetic linkage map construction and identification of Quantitative Trait Loci (QTLs) determining postanthesis drought tolerance and other agronomic traits in bread wheat. PhD Thesis, Halle, Univ., Naturwissenschaftliche Fakultät III
- 288.Liu X-L, Chang X-P, Li R-Z, Jing R-L (2011) Mapping QTLs for seminal root architecture and coleoptile length in wheat. Acta Agronomica Sinica 37(3): 381–388 (RG IF 2016 – 0.12)
- 289.Panigrahi J, Kole P, Kole C (2011) RFLP mapping of loci controlling self-incompatibility in *Brassica campestris* and their comparative mapping with *B. napus* and *B. oleracea*. Biologia Plantarum 55(1):54-60 (IF 2016 – 1.551)
- 290.Ahmadizadeh M, Valizadeh M, Zaefizadeh M, Shahbazi H (2011) Evaluation of interaction between genotype and environments in term of germination and seedling growth in durum wheat landraces. Advances in Environmental Biology 5(4): 551-558
- 291.Wang Z, Mao X, Li A, Chang X, Liu H, Jing R (2011) Functional marker mapping of protein phosphatase 2A structural subunit gene TaPP2Aa in common wheat. Scientia Agricultura Sinica 44 (12): 2411-2421
- 292.Li P, Chen J, Wu P, Zhang J, Chu C, See D, Brown-Guedira G, Zemetra R, Souza E (2011). Quantitative Trait Loci analysis for the effect of Rht-B1 dwarfing gene on coleoptile length and seedling root length and number of bread wheat. Crop Science 51:2561-2568 (IF 2016 – 1.787)
- 293.Yin Zhi-Tong, Meng Fan-Fan, Song Hai-Na et al. (2011) QTL mapping for fast chlorophyll fluorescence parameters in soybean. China Agriculture Science 44(24): 4980-4987
- 294.Isabel Cristina Vinhal-Freitas, Ricardo Franco Tostes, Fernanda Nayara Soares Souza, Jurandir Pereira Segundo, Flávia Rezende Reis (2011). Medidas de emergência e índice de sincronia em sementes de soja osmocondicionadas com polietilenoglicol. Agropecuária Técnica, 32 (1): 102-107
- 295.Cheng K, Xie S, Pu Y, Li S (2012) Comprehensive evaluation on drought resistance in wheat RIL population at seedling stage. Shandong Agricultural Sciences 44: 17-21.
- 296.Rehman Arif MA (2012) Seed longevity and dormancy in wheat (*Triticum aestivum* L.) – phenotypic variation and genetic mapping. PhD Thesis, Martin-Luther-Universität Halle-Wittenberg

- 297.Liu X, Li R, Chang X, Jing R (2013) Mapping QTLs for seedling root traits in a doubled haploid wheat population under different water regimes. *Euphytica* 189(1): 55-61 (IF 2016 – 1.626)
- 298.Jain M, Mittal M, Gadre R (2013) Effect of PEG-6000 Imposed Water Deficit on Chlorophyll Metabolism in Maize Leaves. *Journal of Stress Physiology & Biochemistry* 9 (3): 262-271
- 299.Zhang H, Cui F, Wang L, Li J, Ding A, Zhao C, Bao Y, Yang Q, Wang H (2013) Conditional and unconditional QTL mapping of drought-tolerance-related traits of wheat seedling using two related RIL populations. *Journal of Genetics* 92: 213-231 (IF 2016 – 0.995)
- 300.Khan MA, Iqbal M, Akram M, Ahmad M, Hassan MW, Jamil M (2013) Recent advances in molecular tool development for drought tolerance breeding in cereal crops: a review. *Zemdirbyste-Agriculture* 100 (3):325-334
- 301.Xu Y, Li S, Li L, Zhang H, Xu H, An D (2013) Mapping QTLs for salt tolerance with additive, epistatic and QTL × treatment interaction effects at seedling stage in wheat. *Plant Breeding* 132: 276-283 (IF 2016-1.335)
- 302.Sivakumar R (2013) Studies on physiological, biochemical and molecular basis of drought tolerance in tomato (*Lycopersicon esculentum* Mill.). PhD Thesis, Tamil Nadu Agricultural University Coimbatore, India
- 303.Rebetzke G, Verbyla A, Verbyla K, Morell M, Cavanagh C (2014) Use of a large multiparent wheat mapping population in genomic dissection of coleoptile and seedling growth. *Plant Biotechnology Journal* 12: 219-230 (IF 2016 – 7.443)
- 304.Czyczyłko-Mysza I, Marcińska I, Skrzypek E, Cyganek K, Juzoń K, Karbarz M (2014) QTL mapping for germination of seeds obtained from previous wheat generation under drought. *Central European Journal of Biology* 9: 374-382 (IF 2016 – 0.71)
- 305.Zhang H, Cui F, Wang H (2014) Detection of quantitative trait loci (QTLs) for seedling traits and drought tolerance in wheat using three related recombinant inbred line (RIL) populations. *Euphytica* 196: 313-330 (IF 2016 – 1.626)
- 306.Кърцева Т (2014) Проучване ефекта на мутантни гени върху важни биологични качества при домати (*Solanum lycopersicum* L.) и пшеница (*Triticum aestivum* L.). Дисертация за присъждане на образователната и научна степен “Доктор”, ИФРГ-БАН, София
- 307.Cao P, Ren Y, Zhang K, Teng W, Zhao X, Dong Z, Liu X et al. (2014) Further genetic analysis of a major quantitative trait locus controlling root length and related traits in common wheat. *Molecular Breeding* 33: 975-985 (IF 2016 – 2.465)
- 308.Swain P, Anumalla M, Prusty S, Marndi BC, Rao GJN (2014) Characterization of some Indian native land race rice accessions for drought tolerance at seedling stage. *Australian Journal of Crop Science* 8(3):324-331 (RG IF 2016 – 1.02)
- 309.Li X-M, Chen X-M, Xiao Y-G, Xia X-C, Wang D-S, He Z-H, Wang H-J (2014) Identification of QTLs for seedling vigor in winter wheat. *Euphytica* 198: 199-209 (IF 2016 – 1.626)
- 310.Suárez-González EM, López MG, Délano-Frier JP, Gómez-Leyva JF (2014) Expression of the 1-SST and 1-FFT genes and consequent fructan accumulation in *Agave tequilana* and *A. inaequidens* is differentially induced by diverse (a)biotic-stress related elicitors. *Journal of Plant Physiology* 171: 359-372 (IF 2016 – 3.121)
- 311.Jatoi SA, Latif MM, Arif M, Ahson M, Khan A and Siddiqui SU (2014) Comparative assessment of wheat landraces against polyethylene glycol simulated drought stress. *Science Technology and Development* 33 (1): 1-6
- 312.Ahmad MA, Khan SH, Khan AS, Kazi A-M and Basra SMA (2014) Identification of QTLs for drought tolerance traits on wheat chromosome 2A using association mapping. *International Journal of Agricultural Biology*, 16: 862-870
- 313.Abdolshahi R (2014) Study the inheritance of water use efficiency in bread wheat (*Triticum aestivum*) under drought stress condition. *Electronic Journal of Crop Production* 7(1): 181-199

314. Abdolshahi R, Nazari M, Safarian A, Sadathossini TS, Salarpour M, Amiri H (2015) Integrated selection criteria for drought tolerance in wheat (*Triticum aestivum* L.) breeding programs using discriminant analysis. *Field Crops Research* 174:20-29 (IF 2016 3.048)
315. Acuña-Galindo MA, Mason RE, Subramanian NK, Hays DB (2015) Meta-analysis of wheat QTL regions associated with adaptation to drought and heat stress. *Crop Science* 55:477-492 (IF 2016 – 1.787)
316. Moore C, Rebetzke G (2015) Genomic regions for embryo size and early vigour in multiple wheat (*Triticum aestivum* L.) populations. *Agronomy* 5(2): 152-179 (IF 2016 – 1.614)
317. Khanna-Chopra R, Singh K (2015) Drought resistance in crops: physiological and genetic basis of traits for crop productivity. In: Tripathi BN, Müller M (Eds) *Stress Responses in Plants: Mechanisms of Toxicity and Tolerance* ISBN: 978-3-319-13367-6, pp 267-292
318. Osman Basha P, Sudarsanam G, Madhu Sudhana Reddy M, Siva Sankar N (2015) Effect of PEG induced water stress on germination and seedling development of tomato germplasm. *International Journal of Recent Science Research* 6(5): 4044-4049
319. Tian J, Deng Z, Zhang K, Yu H, Jiang X, Li C (2015) Genetic dissection of stress-tolerance traits in wheat. Chapter 7 in: *Genetic Analyses of Wheat and Molecular Marker-Assisted Breeding*, Volume 1, Springer, jointly published with Science Press, Beijing, pp 445-519
320. Ma J, Du G, Li X, Guo J, Yan G (2016) A QTL on chromosome 3B in bread wheat (*Triticum aestivum*) is associated with leaf width under well-watered and water-deficit conditions. *Plant Molecular Biology Reporter* 34: 690-697 (IF 2016 – 1.932)
321. Qin, Peng, Lin, Yu, Hu, Yaodong, Liu, Kun, Mao, Shuangshuang, Li, Zhanyi, Wang, Jirui, Liu, Yaxi, Wei, Yuming, Zheng, Youliang (2016). Genome-wide association study of drought-related resistance traits in *Aegilops tauschii*. *Genetics and Molecular Biology* 39(3): 398–407 (IF 2015 – 1.341)
322. Ma Y, Kan G, Zhang X, Wang Y, Zhang W, Du H, Yu D (2016) Quantitative Trait Loci (QTL) Mapping for Glycinin and  $\beta$ -Conglycinin Contents in Soybean (*Glycine max* L. Merr.). *Journal of Agriculture and Food Chemistry* 64: 3473–3483 (IF 2016 – 3.154)
323. (ع.م.ر.س.روپ، رفیعی راتخان) (2016) Evaluation of heritability and combining ability of yield and some related traits in bread wheat using half diallel analysis. *Plant Production*, 39(2): 11-26
324. Oyiga BC, Sharma RC, Baum M, Ogbonnaya FC, Léon J, Ballvora A (2017) Allelic variations and differential expressions detected at QTL loci for salt stress tolerance in wheat. *Plant, Cell & Environment*, doi: 10.1111/pce.12898. (IF 2016 – 6.173)
325. Xu, YF, Li, SS, Li, LH et al. (2017) QTL mapping for yield and photosynthetic related traits under different water regimes in wheat. *Molecular Breeding* 37: 34. (IF 2016 – 2.465)
326. Yan Yan, Wang Chunhua, Mu Ping (2017) Genetic analysis of agronomic characters of wheat at seedling stage under drought condition. *Chinese Agricultural Science Bulletin*, 33(3): 8-14
327. Al-Kaaby HK, Abdul-Qadir LH, Abdul-Hussein WF (2017) Role of salicylic acid in the improvement of PEG – induced drought stress in maize (*Zea mays* L.) cv. Baghdad. *Journal of Biology, Agriculture and Healthcare*, 7(2): 25-30
328. Meher, Shivakrishna P, Ashok Reddy K, Manohar Rao D (2017) Effect of PEG-6000 imposed drought stress on RNA content, relative water content (RWC), and chlorophyll content in peanut leaves and roots. *Saudi Journal of Biological Sciences*, <http://dx.doi.org/10.1016/j.sjbs.2017.04.008>

Landjeva S, Korzun V, Stoimenova E, Truberg B, Ganeva G, Börner, A (2008). The contribution of the gibberellin-insensitive semi-dwarfing (*Rht*) genes to genetic variation in wheat seedling growth in response to osmotic stress. *J Agric Sci* 146: 275-286

- 329.Wojciechowski T, Gooding MJ, Ramsay L, Gregory PJ (2009) The effects of dwarfing genes on seedling root growth of wheat. *Journal of Experimental Botany* 60: 2565–2573 (IF 2016 – 5.83)
- 330.Korolev A, Moore T, Saville RJ, Hedden P, Harberd N, Phillips A, Nicholson P and Boulton MI (2009). The role of DELLA proteins in resistance to extreme heat-stress in *Arabidopsis* and cereals. *Plant Temperature Response Networks*, September 23-24, 2009, Durham UK – Abstracts, pp 2-3
- 331.Šíp V, Chrpová J, Žofajová A, Milec Z, Mihalik D, Pánková, K, Snape JW (2011). Evidence of selective changes in winter wheat in middle-European environments reflected by allelic diversity at loci affecting plant height and photoperiodic response. *Journal of Agricultural Science* 149 (3):313-326 (IF 2016 - 1.291)
- 332.González Á, Ayerbe L (2011). Response of coleoptiles to water deficit: growth, turgor maintenance and osmotic adjustment in barley plants (*Hordeum vulgare* L.) *Agricultural Sciences* 2(3):159-166
- 333.Jiang J, Su M, Chen Y, Gao N, Jiao C, Sun Zh, Li F, Wang Ch (2013) Correlation of drought resistance in grass pea (*Lathyrus sativus*) with reactive oxygen species scavenging and osmotic adjustment. *Biologia* 68(2): 231-240 (IF 2016 – 0.759)
- 334.Кърцева Т (2014) Проучване ефекта на мутантни гени върху важни биологични качества при домати (*Solanum lycopersicum* L.) и пшеница (*Triticum aestivum* L.). Дисертация за присъждане на образователната и научна степен “Доктор”, ИФРГ-БАН, София
- 335.Dodig D, Zorić M, Jović M, Kandić V, Stanislavljević R, Šurlan-Momirović G (2015) Wheat seedlings growth response to water deficiency and how it correlates with adult plant tolerance to drought. *Journal of Agricultural Science*, 153: 466-480 (IF 2016 - 1.291)
- 336.Bansal R, Pradheep R, Kumari J, Kumar S, Yadav MC, Gurung B, Kumari NKP, Rana JC (2016) Physiological and biochemical evaluation for drought tolerance in wheat germplasm collected from arid western plains of India. *Indian Journal of Biochemistry & Biophysics* 5: 212-217
- 337.Ayalew H, Liu H, Yan G (2017) Identification and validation of root length QTLs for water stress resistance in hexaploid wheat (*Triticum aestivum* L.) *Euphytica* 213:126. (IF 2016 – 1.626)
- 338.Chen Y, Su YQ, Zhang ChM, Ma J, Mao HT, Yang ZH, et. al. (2017) Comparison of photosynthetic characteristics and antioxidant systems in different wheat strains. *Journal of Plant Growth Regulation* (IF 2016 – 2.073)

**Ganeva G, Petrova T, Law CN, Landjeva S, Sayers L (2008) Plant survival after freezing in wheat ‘Cappelle Desprez’ (‘Bezostaya 1’) intervarietal chromosome substitution lines. Plant Breeding 127 (2), 121–124**

- 339.Eagles HA, Cane K, Trevaskis B (2011) Veery wheats carry an allele of Vrn-A1 that has implications for freezing tolerance in winter wheats. *Plant Breeding* 130 (4): 413-418 (IF 2016- 1.335)

**Nenova V, Merakchiyska M, Ganeva G, Zozikova E and Landjeva S (2009). Physiological responses of wheat (*Triticum aestivum* L.) - *Aegilops sharonensis* introgression lines to excess copper. J Agron Crop Sci 195 (3): 197-203**

- 340.Vertical News: Food Weekly News, 18 June 2009. Online ISSN: 1944-1762
- 341.Величкова, Иrena Богоева (2010) Реакция на житни култури спрямо високи дози мед и инфекция с Fuzarium. Възможности за инактивиране на тежките метали в почвата.- Дисертация за присъждане на образователна и научна степен доктор, Селскостопанска Академия
- 342.Ali S, Zeng F, Qiu L, Zhang G (2011) The effect of chromium and aluminum on growth, root morphology, photosynthetic parameters and transpiration of the two barley cultivars. *Biologia Plantarum* 55: 291-296 (IF 2016 – 1.551)
- 343.Sharma A, Gontia-Mishra I, Srivastava AK (2011) Toxicity of heavy metals on germination and seedling growth of *Salicornia brachiata*. *Journal of Phytology* 3(9): 33-36

344.Dowidar S, Abo-Hamad S, Mohsen A, Khalaf B (2013) Bioremediation of copper stressed *Trigonella foenum graecum*. *Journal of Stress Physiology & Biochemistry*, 9(4): 5-24.

345.Dimitrijević M, Petrović S (2014) Species variation of *Aegilops* genus and heavy metal content in plant habitat soil at southern adriatic localities. *Genetika* 46 (2) 381-400 (RG IF 2016 – 0.44)

346.El Shourbagy MN, Khalil Saad-Allah, SA Foda, EO Razzaky (2017) Impact of Some Halophytic Extracts on the Antioxidant System of Salt-Stressed Safflower (*Carthamus tinctorius L.*). *Journal of Plant Production*, Mansoura Univ. 8 (7): 759 -765

**Kocheva KV, Kartseva T, Landjeva S, Georgiev GI (2009). Physiological response of wheat seedlings to mild and severe osmotic stress. Cereal Res Comm 37 (2): 199-208**

347.Geravandi M, Farshadfar E, Kahrizi, D (2011) Evaluation of some physiological traits as indicators of drought tolerance in bread wheat genotypes. *Russian Journal of Plant Physiology* 58(1): 69-75 (IF 2016 - 0.739)

348.Bencze S, Bamberger Z, Janda T, Balla K, Bedö Z, Veisz O (2011) Drought tolerance in cereals in terms of water retention, photosynthesis and antioxidant enzyme activities. *Central European Journal of Biology* 6:376-387 (IF 2016 – 0.71)

349.Jiang J, Su M, Chen Y, Gao N, Jiao C, Sun Z, Li F, Wang C (2013) Correlation of drought resistance in grass pea (*Lathyrus sativus*) with reactive oxygen species scavenging and osmotic adjustment. *Biologia* 68:231-240 (IF 2016 – 0.759)

350.Farshadfar E, Elyasi P, Hasheminasab H (2013) Incorporation of agronomic and physiological indicators of drought tolerance in a single integrated selection index for screening drought tolerant landraces of bread wheat genotypes. *International Journal of Agronomy and Plant Production* 4: 3314-3325

351.Ortega-Rodes P, Pernus M, Rodés y Eduardo Ortega R (2016) Acclimation of *Nicotiana tabacum* 'Criollo'98' (Solanaceae) to salinity under hydroponic conditions. *Revista Cubana de Ciencias Biológica* 4(3): 16-28

352.Atak M, Mav K (2016) Bazı Serin İklim Tahıllarının İlk Gelişme Döneminde Tuz Stresine Tepkilerinin. *Journal of Agricultural Faculty of Mustafa Kemal University* 21(2):121-129

353.Chen Y, Su YQ, Zhang ChM, Ma J, Mao HT, Yang ZH, et. al. (2017) Comparison of photosynthetic characteristics and antioxidant systems in different wheat strains. *Journal of Plant Growth Regulation* (IF 2016 – 2.073)

**Nagel M, Vogel H, Landjeva S, Buck-Sorlin G, Lohwasser U, Scholz U and Börner A (2009) Seed conservation in ex situ genebanks—genetic studies on longevity in barley. Euphytica 170 (1-2): 5-14**

354.Ulukan H (2011). Plant genetic resources and breeding: current scenario and future prospects. *Int Journal of Agriculture and Biology*, 13: 447–454 (RG IF 2016 – 0.93)

355.Kochanek J, Steadman KJ, Probert RJ, Adkins SW (2011) Parental effects modulate seed longevity: exploring parental and offspring phenotypes to elucidate pre-zygotic environmental influences. *New Phytologist* 191(1):223–233 (IF 2016 – 7.33)

356.Borisjuk L, Rolletschek H, Fuchs J, Melkus G, Neuberger T (2011) Low and High Field Magnetic Resonance for in Vivo Analysis of Seeds. *Materials* 4 (8):1426-1439 (IF 2015 – 2.728)

357.Keller ERJ, Senula A, Zanke C, Grübe M, Kaczmarczyk A (2011) Cryopreservation and in vitro culture - state of the art as conservation strategy for genebanks. *Acta Horticulturae (ISHS)* 918:99-111

358.Van Hintum T, Van Treuren R (2012) Reliability of germination testing of ex situ conserved seeds: A genebank case study on outsourced analyses. *Plant Genetic Resources: Characterisation and Utilisation* 10(2): 134-136 (IF 2016 – 0.612)

359.Scalone R, Albach DC (2012) Degradation of sexual reproduction in *Veronica filiformis* after introduction to Europe. *BMC Evolutionary Biology* 12:233-252 (IF 2016 – 3.221)

- 360.Rehman Arif MA (2012) Seed longevity and dormancy in wheat (*Triticum aestivum* L.) – phenotypic variation and genetic mapping. PhD Thesis, Martin-Luther-Universität Halle-Wittenberg
- 361.de Carvalho MAAP, Bebeli PJ, Bettencourt E, Costa G, Dias S, Dos Santos TMM and Slaski JJ (2013) Cereal landraces genetic resources in worldwide GeneBanks. A review. *Agronomy for Sustainable Development* 33: 177-203 (IF 2016 – 4.101)
- 362.Van Treuren R, Van Hintum T, de Groot EC (2013) Preservation of seed viability during 25 years of storage under standard genebank conditions. *Genetic Resources and Crop Evolution* 60: 1407-1421 (IF 2016 – 1.294)
- 363.Сафина ГФ, Филипенко ГИ (2013) Долговечность семян при хранении и ее прогнозирование методом ускоренного старения. Труды по прикладной ботанике, генетике и селекции. Т. 174. СПб.: ВИР. С. 150, 123-130
- 364.Li G, Na Y-W, Kwon S-W, Park Y-J (2014) Association analysis of seed longevity in rice under conventional and high-temperature germination conditions. *Plant Systematics and Evolution* 300:389-402 (IF 2016 - 1.239)
- 365.Han Z, Ku L, Zhang Z, Zhang J, Guo S, Liu H, Zhang R, Ren Zh, Zhang L, Su H, Dong L, Chen Y (2014) QTLs for Seed Vigor-Related Traits Identified in Maize Seeds Germinated under Artificial Aging Conditions. *PLoS ONE* 9(3): e92535. doi:10.1371/journal.pone.0092535 (IF 2016 – 2.806)
- 366.Dargahi H, Tanya P, Srinivas P (2014) Mapping of the genomic regions controlling seed storability in soybean (*Glycine max* L.) *Journal of Genetics* 93: 365-70 (IF 2016 – 0.995)
- 367.Ku L, Cui X, Cheng F, Guo S, Qi J, Tian Z, Han T, Ren Z, Zhang L, Su H, Chen Y (2014) Genetic dissection of seed vigour under artificial ageing conditions using two joined maize recombinant inbred line populations. *Plant Breeding* 133: 728–737. (IF 2016-1.335)
- 368.Lazar SL, Mira S, Pamfil D, Martinez-Laborde JB (2014) Germination and electrical conductivity tests on artificially aged seed lots of 2 wall-rocket species. *Turkish Journal of Agriculture and Forestry* (2014) 38: 857-864 (IF 2016 – 1.288)
- 369.Groot SPC, de Groot L, Kodde J, van Treuren R (2015) Prolonging the longevity of ex situ conserved seeds by storage under anoxia. *Plant Genetic Resources: Characterization and Utilization* 13: 18-26. (IF 2016 – 0.612)
- 370.Mira S, Estrelles E, González-Benito ME (2015) Effect of water content and temperature on seed longevity of seven Brassicaceae species after 5 years of storage. *Plant Biology* 17: 153-162 (IF 2016 – 2.106)
- 371.Nguyen T-P, Cueff G, Hegedus DD, Rajjou L, Bentsink L (2015) A role for seed storage proteins in *Arabidopsis* seed longevity. *Journal of Experimental Botany* 20: 6399-6413 (IF 2016 – 5.83)
- 372.Ensslin A, Tschoepe O, Burkart M, Joshi J (2015) Fitness decline and adaptation to novel environments in ex-situ plant collections: Current knowledge and future perspectives. *Biological Conservation* 192:394-401 (IF 2016 – 4.022)
- 373.Engels JMM, Šuštar-Vozlič J, Faberová I, Renard M (2015) EUROGENEBANK: Integrated research infrastructure for rational ex situ conservation and use of European plant genetic resources for food and agriculture. DOI: 10.13140/RG.2.1.1377.7769
- 374.Smýkal P, Coyne CJ, Ambrose MJ, Maxted N, Schaefer H, Blair MW, Berger J, Greene SL, Nelson MN, Besharat N, Vymyslický T, Toker C, Saxena RK, Roorkiwal M, Pandey MK, Hu J, Li YH, Wang LX, Guo Y, Qiu LJ, Redden RJ, Varshney RK (2015). Legume Crops Phylogeny and Genetic Diversity for Science and Breeding. *Critical Reviews in Plant Sciences* 34: 43-104 (IF 2016 – 6.825)
- 375.Sasaki K, Takeuchi Y, Miura K, Yamaguchi T, Ando T, Ebitani T, Higashitani A, Yamaya T, Yano M, Sato T (2015) Fine mapping of a major quantitative trait locus, *qLG-9*, that controls seed longevity in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics* 128: 769-778 (IF 2016 – 4.132)

- 376.Fu YB, Ahmed Z, Diederichsen A (2015) Towards a better monitoring of seed ageing under ex situ seed conservation. *Conservation Physiology* 3: doi:10.1093/conphys/cov026. (IF 2016 – 2.323)
- 377.Li Y, Li Y, Yang Q, Zhang J, Zhang J, Qiu L, Wang T (2015) Genomics-based crop germplasm research: advances and perspectives. *Scientia Agricultura Sinica* 48(17): 3333-3353
- 378.Desheva G (2016) The Longevity of Crop Seeds Stored Under Long-term Condition in the National Gene Bank of Bulgaria. *Agriculture (Polnohospodárstvo)*. 62(3): 90-100.
- 379.Listl DM (2017) The geographic scale of genetic variation in common plant species – Implications for genebanks and restoration. PhD, Universität Regensburg, Germany
- 380.Desheva G, Petrova S, Deshev M (2017) Germinability of soybean seeds stored more than 30 years in the Bulgarian national seed genebank. *World Scientific News*, 69: 29-46
- 381.Ahmed Z, Shah ZH, Rehman HM, Shahzad K, Daur I, Elfeel A, Hassan MU, Elsafori AK, Yang SH, Chung G (2017) Genomics: A Hallmark to Monitor Molecular and Biochemical Processes Leading Toward a Better Perceptive of Seed Aging and ex-situ Conservation. *Current Issues of Molecular Biology* 22:89-112 (IF 2016 – 5.25)
- 382.Thormann I, Reeves P, Thumm S, Reiley A, Engels J, Biradar C, ...Richards C (2017) Changes in barley (*Hordeum vulgare* L. subsp. *vulgare*) genetic diversity and structure in Jordan over a period of 31 years. *Plant Genetic Resources*, 1-15. doi:10.1017/S1479262117000028 (IF 2016 – 0.612)
- Ganeva, G., Korzun, V., Landjeva, S., Popova, Z. and Hristov, N. (2010). Genetic diversity assessment of Bulgarian durum wheat (*Triticum durum* Desf.) landraces and modern cultivars using microsatellite markers. *Genetic Resources and Crop Evolution* 57: 273-285 (IF 2008-0.967)**
- 383.Rauf S, da Silva JAT, Khan AA, Naveed A (2010) Consequences of plant breeding on genetic diversity. *International Journal of Plant Breeding* 4: 1-21
- 384.Petrović S (2011). Genetic diversity of winter bread wheat (*Triticum aestivum* L. ssp *vulgare*). PhD Thesis, Josip Juraj Strossmayer University of Osijek, Croatia
- 385.Tsegaye D, Dessalegn T, Dessalegn Y, Share G (2012) Genetic variability, correlation and path analysis in durum wheat germplasm (*Triticum durum* Desf). *Agricultural Research and Reviews* 1(4): 107 – 112
- 386.Tsegaye D, Dessalegn T, Dessalegn Y, Share G (2012) Analysis of genetic diversity in some durum wheat (*Triticum durum* Desf) genotypes grown in Ethiopia. *African Journal of Biotechnology* 11(40): 9606-9611 (RG IF 2016 – 0.44)
- 387.Haile JK, Hammer K, Badebo A, Nachit MM, Röder MS (2013) Genetic diversity assessment of Ethiopian tetraploid wheat landraces and improved durum wheat varieties using microsatellites and markers linked with stem rust resistance. *Genetic Resources and Crop Evolution* 60: 513-527 (IF 2016 – 1.294)
- 388.Nazco R, Peña RJ, Ammar K, Villegas D, Crossa J, Moragues M, Royo C (2013) Variability in glutenin subunit composition of Mediterranean durum wheat germplasm and its relationship with gluten strength. *Journal of Agricultural Science*, 152: 379–393 (IF 2016 – 1.291)
- 389.Todorovska E, Hadjiivanova B, Bozhanova V, Dechev D, Muhovski Y, Panchev I, Abu-Mhadi N, Peycheva V, Ivanova A (2013) Molecular and phenotypic characterization of advanced backcross lines derived from interspecific hybridization of durum wheat. *Biotechnology and Biotechnological Equipment* 27(3): 3760-3771 (IF 2016 – 1.059)
- 390.Gautam A, Sai Prasad SV, Jajoo A, Malviya P (2013) Heritability and correlation of yield and its contributing traits under terminal heat (late sown) situations in durum wheat. *Progressive Research* 8 (2) : 203-208
- 391.Gautam A, Sai Prasad SV, Jajoo A, Malviya P (2013) Stability analysis of yield and its contributing traits in durum wheat under late sown terminal heat stress condition. *Progressive Research* 8 (Special) : 267-272

- 392.Khan MK, Pandey A, Choudhary S, Hakki EE, Akkaya MS, Thomas G (2014) From RFLP to DArT: molecular tools for wheat (*Triticum* spp.) diversity analysis. *Genetic Resources and Crop Evolution* 61(5): 1001-1032 (IF 2016 – 1.294)
- 393.Amallah L, Taghouti M, Rhrib K, Gaboun F, Hassikou R (2014) Genetic variability in agromorphological and quality traits of Mediterranean durum wheat landraces. *Cereal Research Communications* doi: 10.1556/CRC.2014.0029 (IF 2016 – 0.496)
- 394.Леонова ИН, Орловская ОА, Родер МС, Несторов МА, Будашкина ЕБ (2014). Молекулярно-генетическое разнообразие интроверсивных линий мягкой пшеницы (*T. aestivum* /*T. timopheevii*). Вавиловский журнал Генетики и Селекции т. 18, № 4/1: 681-690.
- 395.Li H, Geng S (2015) Assessment of population genetic diversity of *Derris elliptica* (Fabaceae) in China using microsatellite markers. *Industrial Crops and Products* 73: 9-15 (IF 2016 – 3.181)
- 396.Leonova IN, Orlovskaya OA, Röder MS, Nestorov MA, Budashkina EB (2015) Molecular diversity of common wheat introgression lines. (*T. aestivum* /*T. timopheevii*) *Russian Journal of Genetics: Applied Research* 5 (3):191–197
- 397.Liu X, Li Zh, Wang Y, Wang Ch, Zhang H, Ji W (2015). Genetic Diversity Analysis of Aphid-resistant Wheat (*Triticum aestivum*) and Correlation Analysis Between SSR Markers and Resistance to the English Grain Aphid (*Sitobion avenae*). *Journal of Agricultural Biotechnology*, 23(3): 291-301
- 398.Леонова ИН (2015) Генетический контроль устойчивости к грибным болезням у мягкой пшеницы с интроверсиями от *Triticum timopheevii* Zhuk. Диссертация, Институт по цитологии и генетика, СО РАН, Новосибирск
- 399.Wolde T, Eticha F, Alamerew S, Assefa E, Dutamo D (2016) Genetic variability, heritability and genetic advance for yield and yield related traits in durum wheat (*Triticum durum* L.) accessions. *Sky Journal of Agricultural Research* 5(3): 42 – 47
- 400.Phougat D, Godara A, Sethi SK, Sheokand RN (2016) Genetic diversity and association studies for grain yield and its attributing traits in tetraploid wheat (*T. turgidum* subsp. *durum*). *The Bioscan (Supplement on Genetics and Plant Breeding)* 11(3): 2007-2011.

**Landjeva, S., Lohwasser, U. and Börner, A. (2010). Genetic mapping within the wheat D-genome reveals QTL for germination, seed vigour and longevity, and early seedling growth. Euphytica 171 (1): 129-143 (IF 2008-1.403)**

- 401.Ullmannová K, Chloupek O (2011) Influence of genotype and environment on seed vigour of barley. *MendelNet* 176-183
- 402.Wang Q, Cui J, Wang X, Zhang T, Zhou H, Hu T, Han J (2012) Algorithmic models of seed yield and its components in smooth bromegrass (*Bromus inermis* L.) via large sample size under field conditions. *Euphytica*185: 363-375 (IF 2016 – 1.626)
- 403.Kazmi RH, Khan N, Willems LAJ, van Heusden AW, Ligterink W, Hilhorst HWM (2012) Complex genetics controls natural variation among seed quality phenotypes in a recombinant inbred population of an interspecific cross between *Solanum lycopersicum* × *Solanum pimpinellifolium*. *Plant, Cell & Environment* 35(5): 929-951 (IF 2016 – 6.173)
- 404.Paliwal R, Röder MS, Kumar U, Srivastava JP, Joshi AK (2012) QTL mapping of terminal heat tolerance in hexaploid wheat (*T. aestivum* L.). *Theoretical and Applied Genetics* 125: 561-575 (IF 2016 – 4.132)
- 405.Thu-Phuong Nguyen, Paul Keizer, Fred van Eeuwijk, Sjef Smeekens, Leónie Bentsink (2012) Natural variation for seed longevity and seed dormancy are negatively correlated in *Arabidopsis thaliana*. *Plant Physiology* 160: 2083-92 (IF 2015 – 6.280)

- 406.Bogard M (2012) Analyse génétique et écophysiologique de l'écart à la relation teneur en protéines - rendement en grains chez le blé tendre (*Triticum aestivum* L.). PhD Thesis, Université Blaise Pascal, Clermont-Ferrand II, France
- 407.Rehman Arif MA (2012) Seed longevity and dormancy in wheat (*Triticum aestivum* L.) – phenotypic variation and genetic mapping. PhD Thesis, Martin-Luther-Universität Halle-Wittenberg
- 408.Kumar A, Simons K, Iqbal MJ, Jiménez MM, Bassi FM et al. (2013) Physical mapping resources for large plant genomes: radiation hybrids for wheat D-genome progenitor *Aegilops tauschii*. BMC Genomics, 13:597 doi:10.1186/1471-2164-13-597 (IF 2016 – 3.729)
- 409.Zhang H, Cui F, Wang L, Li J, Ding A, Zhao C, Bao Y, Yang Q, Wang H (2013) Conditional and unconditional QTL mapping of drought-tolerance-related traits of wheat seedling using two related RIL populations. Journal of Genetics 92(2): 213-231 (IF 2016 – 0.995)
- 410.Kazmi RH (2013) Genes for seed quality: Integrating physiology and genetical genomics to mine for seed quality genes in tomato. PhD Thesis, Wageningen University, Wageningen, NL ISBN 978-94-6173-520-1
- 411.Czyczyło-Mysza I, Marcińska I, Skrzypek E, Cyganek K, Juzoń K, Karbarz M (2014) QTL mapping for germination of seeds obtained from previous wheat generation under drought. Central European Journal of Biology 9: 374-382 (IF 2016 – 0.71)
- 412.Zhang H, Cui F, Wang H (2014) Detection of quantitative trait loci (QTLs) for seedling traits and drought tolerance in wheat using three related recombinant inbred line (RIL) populations. Euphytica 196: 313-330 (IF 2016 – 1.626)
- 413.Xie L, Tan Zh, Zhou Y, Xu R, Feng L, Xing Y, Qi X (2014) Identification and fine-mapping of quantitative trait loci for seed vigor in germination and seedling establishment in rice. Journal of Integrative Plant Biology 56(8): 749-59 (IF 2016 – 3.962)
- 414.Li X-M, Chen X-M, Xiao Y-G, Xia X-C, Wang D-S, He Z-H, Wang H-J (2014) Identification of QTLs for seedling vigor in winter wheat. Euphytica 198(2): 199-209 (IF 2016 – 1.626)
- 415.Engelmann U (2014) Kartierung und züchterische Nutzung von Resistzenzen gegen die Weizenblattdürre (*Pyrenophora tritici-repentis*). PhD Thesis, Julius Kühn-Institut (JKI), Quedlinburg, Germany
- 416.Mahmood T, Trethowan R (2015) Crop Breeding for Conservation Agriculture. In: Farooq M, Siddique K (Eds). Conservation Agriculture, pp 159-179. Springer International Publishing
- 417.Sasaki K, Takeuchi Y, Miura K, Yamaguchi T, Ando T, Ebitani T, Higashitani A, Yamaya T, Yano M, Sato T (2015) Fine mapping of a major quantitative trait locus, *qLG-9*, that controls seed longevity in rice (*Oryza sativa* L.). Theoretical and Applied Genetics 128: 769-778 (IF 2016 – 4.132)
- 418.Nguyen T-P, Cueff G, Hegedus DD, Rajjou L, Bentsink L (2015) A role for seed storage proteins in *Arabidopsis* seed longevity. Journal of Experimental Botany 66:6399-413 (IF 2016 – 5.83)
- 419.Kammili A, Jatohu J (2015) Differential longevity of castor (*Ricinus communis* L.) germplasm conserved under uncontrolled storage conditions across extended periods, Journal of Crop Improvement, 29, 6, 706-719
- 420.Tian J, Deng Z, Zhang K, Yu H, Jiang X, Li C (2015) Genetic analysis of main physiological and morphological traits. Chapter 6 in: Genetic Analyses of Wheat and Molecular Marker-Assisted Breeding, Volume 1, Springer, jointly published with Science Press, Beijing, pp 351-443
- 421.Sano N, Rajjou L, North HM, Debeaujon I, Annie Marion-Poll AM, Mitsunori Seo M (2016). Staying Alive: Molecular Aspects of Seed Longevity. Plant and Cell Physiology 57(4): 660–674 (IF 2016 – 4.76)
- 422.Lima JJP (2016) Physiological and molecular studies during acquisition of longevity in soybean (*Glycine max* (L.) Merrill) seeds. PhD Thesis, Universidade Estadual Paulista “Julio de Mesquita Filho”, Botucatu

- 423.Sharma D, Singh R, Rane J, Gupta VK, Mamrutha HM, Tiwari R (2016) Mapping quantitative trait loci associated with grain filling duration and grain number under terminal heat stress in bread wheat (*Triticum aestivum* L.). *Plant Breeding*, 135 (5): 538-545 (IF 2016-1.335)
- 424.Boudiar R, Casas AM, Cantalapiedra CP, Pilar Gracia M, Igartua E (2016) Identification of quantitative trait loci for agronomic traits contributed by a barley (*Hordeum vulgare*) Mediterranean landrace. *Crop and Pasture Science* 67(1): 37-46 (IF 2016 – 1.804)
- 425.Wang B, Zhang Zh, Fu Zh, Liu Z, Hu Y, Tang J (2016) Comparative QTL analysis of maize seed artificial aging between an immortalized F2 population and its corresponding RILs. *The Crop Journal*, 4(1):30-39
- 426.Saisho D, Takumi S, Matsuoka Y (2016) Salt tolerance during germination and seedling growth of wild wheat Aegilops tauschii and its impact on the species range expansion. *Scientific Reports* 6, Article No 38554, doi: 10.1038/srep38554 (IF 2016 – 4.259)
- 427.Min Tan, Fang Liao, Lintao Hou, Jia Wang, Lijuan Wei, Hongju Jian, Xinfu Xu, Jiana Li, Liezhao Liu (2017) Genome-wide association analysis of seed germination percentage and germination index in *Brassica napus* L. under salt and drought stresses. *Euphytica* 213:40 (IF 2016 – 1.626)
- 428.Jaradat AA (2017) Phenotypic and Ionomics Profiling of *Triticum aestivum* × *Aegilops tauschii* Introgression Lines. *Crop Science* 0. doi:10.2135/cropsci2016.09.0797 (IF 2016 – 1.787)
- 429.Arooj F, Qayyum A, Mahmood S (2017) Responses of wheat seedling to varying moisture conditions and relationship between morphological and molecular characterization. *International Journal of Environmental & Agriculture Research* 3(4): 124-135
- 430.Chen GF, Wu RG, Li DM, Yu HX, Deng Zh,Tian JC (2017) Genomewide association study for seedling emergence and tiller number using SNP markers in an elite winter wheat population. *Journal of Genetics* 96: 177–186 (IF 2016 – 0.995)
- 431.Silva TA (2017) Influence of residual lime in the acquisition of the physiological quality of wheat seeds produced in direct sowing system. PhD Thesis, Universidade Estadual Paulista “Júlio Mesquita Filho” Campus de Botucatu, Brazil
- 432.Adu-Gyamfi PKK (2017) The physiological basis of drought tolerance identified from genetic association analysis in wheat (*Triticum aestivum* L.). PhD Thesis, The University of Sydney, Australia

**Melnikova NV, Ganeva GD, Popova ZG, Landjeva SP, Kudryavtsev AM (2010) Gliadins of Bulgarian durum wheat (*Triticum durum* Desf.) landraces: genetic diversity and geographical distribution Genetic Resources and Crop Evolution 57:587–595**

- 433.Nazco R, Villegas D, Ammar K, Peña R,J, Moragues M, Royo C (2012) Can Mediterranean durum wheat landraces contribute to improved grain quality attributes in modern cultivars? *Euphytica* 185:1-17 (IF 2016 – 1.626)
- 434.Ruiz M, Giraldo P, Royo C, Villegas D, Jose Aranzana M, Carrillo JM (2012) Diversity and genetic structure of a collection of spanish durum wheat landraces. *Crop Science* 52(5):2262-2275 (IF 2016 – 1.787)
- 435.Soriano JM, Villegas D, Aranzana MJ, García del Moral LF, Royo C (2016) Genetic structure of modern durum wheat cultivars and mediterranean landraces matches with their agronomic performance. *PLoS ONE* 11(8): e0160983. doi:10.1371/journal.pone.0160983 (IF 2016 – 2.806)
- 436.Utebayev M, Dashkevich S, Babkenov A, Shtefan G, Fahrudenova I, Bayahmetova S, Sharipova B, Kaskarbayev Z, Shavrukov Y (2016) Application of gliadin polymorphism for pedigree analysis in common wheat (*Triticum aestivum* L.) from Northern Kazakhstan. *Acta Physiol Plant* 38:204

**Kolev S, Ganeva G, Christov N, Belchev I, Kostov K, Tsenov N, Rachovska G, Landjeva S, Ivanov M, Abu-Mhadi N, Todorovska E (2010). Allele variation in loci for adaptive response and plant height and its effect on grain yield in wheat. Biotechnology & Biotechnological Equipment. 24 (2): 1807-1813; ISSN 1310-2818. (IF 2009 0.291)**

- 437.Vinod Sharma JB, Tomar RS, Singh B, Tomar SMS (2012) Breeding for earliness and thermo-tolerance in wheat suited to early, late and very late sowing in northwestern India. Indian Journal of Genetics and Plant Breeding 72: 15-24
- 438.Hu W-M, Kan H-H, Wang W and Xu C-W (2013) Statistical genetics approach for functional difference identification of allelic variations and its application. Acta Agronomica Sinica 11 (RG IF 2016 – 0.12)
- 439.Markova L, Tomchev N (2013) Seasonal dynamics of cereal aphids and predators on various wheat genotypes (*Triticum aestivum* L.). In: Proc 1st Global Virtual Conference, April, 8. - 12. 2013, pp. 669- 673, <http://www.gv-conference.com>
- 440.Кърцева Т (2014) Проучване ефекта на мутантни гени върху важни биологични качества при домати (*Solanum lycopersicum* L.) и пшеница (*Triticum aestivum* L.). Дисертация за присъждане на образователната и научна степен “Доктор”, ИФРГ-БАН, София
- 441.Akman H, Bruckner P (2016) Marker assisted selection for photoperiod insensitive Ppd-D1a allele in winter wheat breeding program. International Journal of Ecosystems and Ecology Sciences 6 (2): 147-150 (IF 2016 1.463)
- 442.Khumalo TP, Barnard A, Maphosa L, Tsilo TJ (2017) Impact of Growth Habit and Architecture Genes on Adaptation and Performance of Bread Wheat. Chapter 5, In:Wanyera R and Owuiche J (Eds) Wheat Improvement, Management and Utilization, InTech, pp. 77-95

**Kocheva K, Kartseva T, Landjeva S, Georgiev G (2010). Parameters of cell membrane stability and levels of oxidative stress in leaves of wheat seedlings treated with PEG 6000. General and Applied Plant Physiology 35(3-4): 127-133.**

- 443.Zala HN, Bosamia TC, Kulkarni KS, Shukla YM (2014) Physiological and biochemical characterization for drought stress at seedling stage in wheat genotypes. International Journal of Bio-resource and Stress Management 5(3):386-394
- 444.Anatala Tushar J, Gajera HP, Mandavia MK, DaveRajesh A, Kothari Vishal V, Golakiya BA (2015) Leaf proteome alterations in tolerant pearl millet (*Pennisetum glaucum* L.) genotype under water stress. International Journal of Agriculture, Environment and Biotechnology 8(3): 539-549 (IF 2015 – 0.546)
- 445.Öztür A, Taşkesenligil B, Haliloglu K, Aydin M, Çağlar Ö (2016) Evaluation of bread wheat genotypes for early drought resistance via germination under osmotic stress, cell membrane damage, and paraquat tolerance. Turkish Journal of Agriculture and Forestry, 40: 146-159 (IF 2016 – 1.288)

**Landjeva S, Karceva T, Korzun V, Ganeva G (2012). Seedling growth under osmotic stress and agronomic traits in Bulgarian semi-dwarf wheat – comparison of genotypes with Rht8 and/or Rht-B1 genes. Crop and Pasture Science, 62(12): 1017-1025 (IF 2010 1.280)**

- 446.Чеботар ГО (2012) Алелі генів короткостебловості Rht8, Rht-B1, Rht-D1, нечутливості до фотоперіоду Ppd-D1 м'якої пшениці та їх ефекти на агрономічні ознаки // Дисертація на здобуття наукового ступеня канд. біол. наук. Одеса, 185 с
- 447.Divashuk MG, Bespalova LA, Vasiliyev AV, Fesenko IA, Puzyrnaya OYu, Karlov GI (2013) Reduced height genes and their importance in winter wheat cultivars grown in southern Russia. Euphytica 190: 137-144 (IF 2016 – 1.626)
- 448.Mohan A, Schillinger WF, Gill KS (2013) Wheat seedling emergence from deep planting depths and its relationship with coleoptile length. PLoS ONE 8(9), e73314. doi:10.1371/journal.pone.0073314 (IF 2016 – 2.806)
- 449.Tsenov N, Gubatov T, Tsenova E (2015) Effect of height of stem on the productivity of winter common wheat. Agricultural Science and Technology 7(2): 179-185

450.Chebotar GA, Chebotar SV, Motsnyy II (2016) Pleiotropic effects of gibberellin-sensitive and gibberellin-insensitive dwarfing genes in bread wheat of the southern step region of the Black Sea. Cytology and Genetics 50 (1): 20-27. (IF 2016 – 0.324)

451.Гончарова AI, Моцний II, Чеботар СВ (2016) Вплив алелів генів короткостебловості на довжину колеоптиля пшениці м'якої в умовах осмотичного стресу. Вісник Одеського національного університету. Біологія. 21, 1(38): 89-99

**Landjeva S, K Kocheva, T Karceva, A Seps, I Molnár, A Schneider, G Ganeva, G Georgiev, M Molnár-Láng (2012). Molecular cytogenetic identification of a wheat-Aegilops geniculata Roth spontaneous chromosome substitution and its effects on the growth and physiological responses of seedlings to osmotic stress. Plant Breeding 131(1): 81-87 (IF 2010 1.391)**

452.Zhao Hui, Zhang Wei, Wang Jing, Li Fu-Rong, Ji Jun, Li Jun-Ming (2013) Research progress on improving wheat drought tolerance via exotic gene introduction. Chinese Journal of Eco-Agriculture 1(3): 267–273

453.Zeinali G, Mirzaghaderei G, Badakhshan H (2013) Genome structure and salt stress response of some segregated lines from wheat and Tritipyrum crosses. Cytologia 78(4): 1–11 (IF 2016 – 0.913)

**Ganeva G, T Petrova, S Landjeva, E Todorovska, S Kolev, G Galiba, F Szira, A Bálint (2013) Frost tolerance in winter wheat cultivars: different effects of chromosome 5A and association with microsatellite alleles. Biologia Plantarum 57 (1): 184-188**

454.Zheng B, Chapman SC, Christopher JY, Frederiks TM, Chenu K (2015) Frost trends and their estimated impact on yield in the Australian wheatbelt. Journal of Experimental Botany 66 (12) doi:10.1093/jxb/erv163 (IF 2016 – 5.83)

455.Afzal F, Chaudhari SH, Gul A, Farooq A, Ali H, Nisar S, Sarfraz B, Shehzadi KJ, Mujeeb-Kazi A. (2015) Bread wheat (*Triticum aestivum L.*) under biotic and abiotic stresses. In: Hakeem K R (Ed) Crop Production and Global Environmental Issues, Springer, ISBN 978-3-319-23-161-7, pp 293-318

**Kocheva KV, Landjeva SP and Georgiev GI (2014) Variation in ion leakage parameters of two wheat genotypes with different Rht-B1 alleles in response to drought. Journal of Biosciences 39: 753-759 DOI 10.1007/s12038-014-9471-7 ISSN: 0973-7138 (IF 2013-1.939)**

456.Barakat MN, Al-Doss AA, Moustafa KA, Elshafei AA, Salem AK (2015) Identification of QTLs for four physiological traits in an advanced backcross population of wheat under drought stress. Plant OMICS 8(2):122-129 (IF 2016 - 0.794)

**Kocheva K, Nenova V, Karceva T, Petrov P, Georgiev GI, Börner A, Landjeva S (2014) Changes in water status, membrane stability and antioxidant capacity of wheat seedlings carrying different Rht-B1 dwarfing alleles under drought stress. J Agro Crop Sci 200 (2): 83–91**

457.Munjal R, Dhanda S S (2016) Assessment of drought resistance in Indian wheat cultivars for morpho-physiological traits. Ekin Journal 2(1):74-81

458.Yan J, Zhang N, Wang N, LI Y, Zhang S, Wang S (2016) Variations in adaptation strategies of wheat cultivar replacements under short-term osmotic stress. Pakistan Journal of Botany, 48(3): 917-924 (IF 2016 – 0.658)

459.Jiakun YAN, Suiqi ZHANG. Effects of dwarfing genes on water use efficiency of bread wheat[J]. Front. Agr. Sci. Eng., 2017, 4(2): 126-134.

**Ganeva G, Landjeva S, Belchev I, Koleva L (2014) Characterization of two wheat doubled haploid populations for resistance to common bunt and its association with agronomic traits. Cereal Res communications 42(3): 484–494**

460.Lantos C, Pauk J (2016) Anther culture as an effective tool in winter wheat (*Triticum aestivum* L.) breeding. Russian Journal of Genetics 52(8): 794–801 (IF 2016 – 0.55)

**Börner A, Landjeva S, Nagel M, Rehman Arif MA, Allam M, Agacka M, Doroszewska T, Lohwasser U (2014) Plant Genetic Resources for Food and Agriculture (PGRFA) – Maintenance and Research. Genetics and Plant Physiology 4: 13–21**

461.Listl DM (2017) The geographic scale of genetic variation in common plant species – Implications for genebanks and restoration. PhD, Universität Regensburg, Germany

**Landjeva S, Ganeva G, Korzun V, Palejev D, Chebotar S, Kudrjavtsev A (2015) Genetic diversity of old bread wheat germplasm from the Black Sea region evaluated by microsatellites and agronomic traits. Plant Genetic Resources: Characterization and Utilization 13 (2): 119-130**

462.Kolesnyk O, Khokhlov A (2015) Association analysis of TKW in Ukrainian bread wheat varieties using SSR markers. Modern Science 1: 150-158

463.Brbaklić L. (2015) Molecular and association analysis of spike - related traits in wheat (*Triticum aestivum* L.). PhD Thesis, University of Belgrade

464.Oscar G, Blangiforti S, Venora G (2017) Wheat landraces identification through glumes image analysis. Computers and Electronics in Agriculture 141: 223-231 (IF 2016 – 2.201)

София, декември 2017

Изготвил:

(доц. д-р Светлана Мишева)